

## Empirical Process Proof of the Asymptotic Distribution of Sample Quantiles

**Definition:** Given  $\theta \in (0, 1)$ , the  $\theta^{\text{th}}$  **quantile** of a random variable  $\tilde{X}$  with CDF  $F$  is defined by:

$$\mu_\theta \equiv F^{-1}(\theta) = \inf\{x \mid F(x) \geq \theta\}.$$

Note that  $\mu_{.5}$  is the *median*,  $\mu_{.25}$  is the 25<sup>th</sup> percentile, etc. Further if we define the 0<sup>th</sup> quantile as  $\mu_0 = \lim_{\theta \rightarrow 0} \mu_\theta$  and define  $\mu_1$  similarly, it is easy to see that these are the lower and upper points in the support of  $\tilde{X}$  (i.e. the minimum and maximum possible values of  $\tilde{X}$  which might be  $-\infty$  and  $+\infty$  if  $\tilde{X}$  has unbounded support). Note also that if  $F$  is strictly increasing in a neighborhood of  $\mu_\theta$ , then  $\mu_\theta = F^{-1}(\theta)$  is the usual inverse of the CDF  $F$ . If  $F$  happens to have “flat” sections, say an interval of points  $x$  satisfying  $F(x) = \theta$ , then  $\mu_\theta$  is the smallest  $x$  in this interval. The following lemma, a slightly modified version of a lemma from R. J. Serfling, (1980) *Approximation Theorems of Mathematical Statistics* Wiley, New York, provides some basic properties of the *quantile function*  $F^{-1}(\theta)$ :

**Lemma 1:** Let  $F$  be a CDF. The quantile function  $F^{-1}(\theta)$ ,  $\theta \in (0, 1)$  is non-decreasing and left continuous, and satisfies:

1.  $F^{-1}(F(x)) \leq x$ ,  $-\infty < x < \infty$
2.  $F(F^{-1}(\theta)) \geq \theta$ ,  $0 < \theta < 1$
3. If  $F$  is strictly increasing in a neighborhood of  $\mu_\theta = F^{-1}(\theta)$  we have:  $F(F^{-1}(\theta)) = \theta$  and  $F^{-1}(F(\mu_\theta)) = \mu_\theta$ .
4.  $F(x) \geq \theta$  if and only if  $x \geq F^{-1}(\theta)$ .

**Definition:** Let  $(\tilde{X}_1, \dots, \tilde{X}_N)$  be a random sample of size  $N$  from a CDF  $F$ . Then the **sample quantile**  $\hat{\mu}_\theta$ ,  $\theta \in (0, 1)$  is defined by:

$$\hat{\mu}_\theta = F_N^{-1}(\theta),$$

where  $F_N$  is the empirical CDF defined by:

$$F_N(x) = \frac{1}{N} \sum_{i=1}^N I\{\tilde{X}_i \leq x\}.$$

Thus  $\hat{\mu}_{.5} = F_N^{-1}(.5)$  is the *sample median*  $F_N^{-1}(.5) = \text{med}(\tilde{X}_1, \dots, \tilde{X}_N)$ , and  $\hat{\mu}_0 = F_N^{-1}(0)$  is the *sample minimum*,  $F_N^{-1}(0) = \min(\tilde{X}_1, \dots, \tilde{X}_N)$ , and  $\hat{\mu}_1 = F_N^{-1}(1)$  is the *sample maximum*,  $F_N^{-1}(1) = \max(\tilde{X}_1, \dots, \tilde{X}_N)$ . Since empirical CDF's have jumps of size  $1/N$  (unless more than one of the  $\{\tilde{X}_i\}$ 's take the same value), then we can bound the maximum difference between  $\theta$  and  $F(F^{-1}(\theta))$  in Lemma 1-2 as follows:

**Lemma 2:** Let  $(\tilde{X}_1, \dots, \tilde{X}_N)$  be a random sample from a CDF  $F$  and suppose that in this sample each  $\tilde{X}_i$  happens to be distinct, so that by reindexing we have  $\tilde{X}_1 < \tilde{X}_2 < \dots < \tilde{X}_{N-1} < \tilde{X}_N$ . Then for all  $\theta \in (0, 1)$  we have:

$$|F_N(F_N^{-1}(\theta)) - \theta| \leq \frac{1}{N}.$$

The following theorem shows that the asymptotic distribution of the sample quantiles  $\hat{\mu}_\theta$  for  $\theta \in (0, 1)$  are normally distributed. It is important to note that we exclude the two cases  $\theta = 0$  and  $\theta = 1$  in this theorem since the asymptotic distribution of these *extreme value statistics* is very different and generally non-normal.

**Theorem:** Let  $(\tilde{X}_1, \dots, \tilde{X}_N)$  be IID draws from a CDF  $F$  with continuous density  $f$ . Then if  $f(\mu_\theta) > 0$ , we have:

$$\sqrt{N}(\hat{\mu}_\theta - \mu_\theta) = \sqrt{N}(F_N^{-1}(\theta) - F^{-1}(\theta)) \implies N(0, \sigma^2),$$

where:

$$\sigma^2 = \frac{\theta(1-\theta)}{f(\mu_\theta)^2}.$$

**Proof:** The Central Limit Theorem for IID random variables implies that for any  $x$  in the support of  $F$  we have:

$$\sqrt{N}(F_N(x) - F(x)) \implies N(0, \gamma^2),$$

where  $\gamma^2 = F(x)[1 - F(x)]$ . Letting  $x = \mu_\theta = F^{-1}(\theta)$  and using Lemma 1-3 we have:

$$\sqrt{N}(F_N(\mu_\theta) - F(\mu_\theta)) = \sqrt{N}(F_N(F^{-1}(\theta)) - F(F^{-1}(\theta))) \implies N(0, \theta(1-\theta)).$$

Furthermore, the property of *stochastic equicontinuity* from the theory of *empirical processes* (see D. Andrews, (1996) *Handbook of Econometrics* (vol. 4) for an accessible introduction and definition of stochastic equicontinuity), we have that the result given above is unaffected if we replace  $\mu_\theta$  by a consistent estimate  $\hat{\mu}_\theta$ :

$$\sqrt{N}(F_N(\hat{\mu}_\theta) - F(\hat{\mu}_\theta)) = \sqrt{N}(F_N(F_N^{-1}(\theta)) - F(F_N^{-1}(\theta))) \implies N(0, \theta(1-\theta)).$$

Now note that Lemma 1-2 implies that

$$\sqrt{N}(F_N(F_N^{-1}(\theta)) - F(F_N^{-1}(\theta))) \geq \sqrt{N}(\theta - F(F_N^{-1}(\theta))).$$

However since the true CDF  $F$  has a density, the probability of observing duplicate  $\{\tilde{X}_i\}$ 's is zero, so Lemma 2 implies that with probability 1 we have:

$$\sqrt{N}(F_N(F_N^{-1}(\theta)) - F(F_N^{-1}(\theta))) = \sqrt{N}(\theta - F(F_N^{-1}(\theta))) + O_p(1/\sqrt{N}),$$

which implies that:

$$\sqrt{N}(\theta - F(F_N^{-1}(\theta))) \implies N(0, \theta(1-\theta)).$$

Now we apply the Delta theorem, i.e. we do a Taylor series expansion of  $F(F_N^{-1}(\theta))$  about the limiting point  $\mu_\theta = F^{-1}(\theta)$  to get:

$$F(F_N^{-1}(\theta)) = F(F^{-1}(\theta)) + f(\tilde{\mu}_\theta) [F_N^{-1}(\theta) - F^{-1}(\theta)],$$

where  $\tilde{\mu}_\theta$  is a point on the line segment between  $\hat{\mu}_\theta$  and  $\mu_\theta$ . Using the result above and Lemma 1-3 we have:

$$\sqrt{N} \left( F_N^{-1}(\theta) - F^{-1}(\theta) \right) = \sqrt{N} \left( \frac{\theta - F(F_N^{-1}(\theta))}{-f(\tilde{\mu}_\theta)} \right) \implies N(0, \sigma^2),$$

where we have used Slutsky's Theorem and the fact that  $\tilde{\mu}_\theta \rightarrow \mu_\theta$  since  $\hat{\mu}_\theta \rightarrow \mu_\theta$  with probability 1.