

LIMIT DISTRIBUTION THEORY FOR MAXIMUM LIKELIHOOD ESTIMATION OF A LOG-CONCAVE DENSITY

BY FADOUA BALABDAOUI, KASPAR RUFIBACH¹ AND JON A. WELLNER²

Universite Paris-Dauphine and University of Göttingen, University of Zurich and University of Washington

We find limiting distributions of the nonparametric maximum likelihood estimator (MLE) of a log-concave density, that is, a density of the form $f_0 = \exp \varphi_0$ where φ_0 is a concave function on \mathbb{R} . The pointwise limiting distributions depend on the second and third derivatives at 0 of H_k , the “lower envelope” of an integrated Brownian motion process minus a drift term depending on the number of vanishing derivatives of $\varphi_0 = \log f_0$ at the point of interest. We also establish the limiting distribution of the resulting estimator of the mode $M(f_0)$ and establish a new local asymptotic minimax lower bound which shows the optimality of our mode estimator in terms of both rate of convergence and dependence of constants on population values.

1. Introduction.

1.1. *Log-concave densities.* A probability density f on the real line is called log-concave if it can be written as

$$f(x) = \exp \varphi(x)$$

for some concave function $\varphi: \mathbb{R} \rightarrow [-\infty, \infty)$. We let \mathcal{LC} denote the class of all log-concave densities on \mathbb{R} . As shown by Ibragimov (1956), a density function f is log-concave if and only if its convolution with any unimodal density is again unimodal. Thus, the class of log-concave densities is often referred to as the class of “strongly unimodal” densities. Furthermore, the class \mathcal{LC} of log-concave densities is exactly the class of Polyá frequency functions of order 2, PFF_2 as noted by Pal, Woodroffe and Meyer (2007); see also Dharmadhikari and Joag-Dev (1988), page 150, and Marshall and Olkin (1979), page 492.

The log-concave shape constraint is appealing for many reasons:

(1) Many parametric models, for a certain range of their parameters, are in fact log-concave, for example, normal, uniform, gamma(r, λ) for $r \geq 1$, beta(a, b) for

Received January 2008.

¹Supported in part by the Swiss National Science Foundation.

²Supported in part by NSF Grant DMS-05-03822 and NI-AID Grant 2R01 AI291968-04.
AMS 2000 subject classifications. Primary 62N01, 62G20; secondary 62G05.

Key words and phrases. Asymptotic distribution, integral of Brownian motion, envelope process, log-concave density estimation, lower bounds, maximum likelihood, mode estimation, nonparametric estimation, qualitative assumptions, shape constraints, strongly unimodal, unimodal.

$a \geq 1$ and $b \geq 1$, generalized Pareto, Gumbel, Fréchet, logistic or Laplace, to mention only some of these models. Therefore, assuming log-concavity offers a flexible nonparametric alternative to purely parametric models. Note that a log-concave density need not be symmetric.

(2) Every log-concave density is automatically unimodal. Furthermore, log-concavity of a density f immediately implies specific shape constraints for certain functions derived from f [see Barlow and Proschan (1975), Marshall and Olkin (1979, 2007), Dharmadhikari and Joag-Dev (1988), An (1998) and Bagnoli and Bergstrom (2005)]. Thus, having an estimator (and its limiting distribution) for f at hand provides, almost automatically, estimators (and limiting distributions) for those functions. Corollary 2.3 illustrates this for the hazard rate.

(3) Although the nonparametric MLE of a unimodal density does not exist [see, e.g., Birgé (1997)], the nonparametric MLE of a log-concave density exists, is unique and has desirable consistency and rates of convergence properties. Thus, the class of log-concave (or strongly unimodal) densities may be a useful and valuable surrogate for the larger class \mathcal{U} of unimodal densities.

(4) Tests for multimodality and mixing can be based on a semiparametric model with densities of the form $f_{c,\varphi}(x) = \exp(\varphi(x) + cx^2)$, where φ is concave and $c > 0$, as shown by Walther (2002).

(5) Chang and Walther (2007) further show that the EM-algorithm can be extended to work for log-concave component densities.

(6) First attempts to estimate a log-concave density in \mathbb{R}^d were made by Cule, Gramacy and Samworth (2007).

(7) The log-concave density estimator can be used to improve accuracy in the estimation of the so-called “tail index” of a generalized Pareto distribution [see Müller and Rufibach (2009)].

(8) It should be noted that no arbitrary choices such as bandwidth, kernel or prior are involved in the estimation of a log-concave density; these are all obviated by this shape restriction.

(9) We expect good adaptivity properties of the MLE \hat{f}_n in the class \mathcal{LC} .

For properties of (random variables with) log-concave densities, we refer to Dharmadhikari and Joag-Dev (1988), Marshall and Olkin (1979) and Rufibach (2006). Log-concavity of a density f implies certain shape constraints for functions derived from f , such as the distribution function, the tail or hazard function. See An (1998) for comparisons with the related notion of a log-convex density.

1.2. Log-concave density estimation. Now let $X_{(1)} < X_{(2)} < \dots < X_{(n)}$ be the order statistics of n independent random variables X_1, \dots, X_n , distributed according to a log-concave probability density $f_0 = \exp \varphi_0$ on \mathbb{R} . The distribution function corresponding to f_0 is denoted by F_0 .

The maximum likelihood estimator (MLE) of a log-concave density was introduced in Rufibach (2006) and Dümbgen and Rufibach (2009). Algorithmic aspects were treated in Rufibach (2007) and in a more general framework in Dümbgen, Hüsler and Rufibach (2007), while consistency with respect to the Hellinger metric was established by Pal, Woodroffe and Meyer (2007), and rates of convergence of \hat{f}_n and \hat{F}_n were established by Dümbgen and Rufibach (2009). Since the derivation of the MLE of a log-concave density is extensively treated in these references, we only briefly recall its definition and the properties relevant for this paper.

If \mathcal{C} denotes the class of all concave functions $\varphi : \mathbb{R} \rightarrow [-\infty, \infty)$, the estimator $\hat{\varphi}_n$ of φ_0 is the maximizer of the “adjusted” criterion function

$$L(\varphi) = \int_{\mathbb{R}} \varphi(x) d\mathbb{F}_n(x) - \int_{\mathbb{R}} \exp \varphi(x) dx$$

over \mathcal{C} , where \mathbb{F}_n is the empirical distribution function of the observations. The log-concave density estimator is then $\hat{f}_n := \exp \hat{\varphi}_n$, which exists and is unique.

1.3. *Characterization of $\hat{\varphi}_n$.* For any continuous piecewise linear function $h_n : [X_{(1)}, X_{(n)}] \rightarrow \mathbb{R}$, such that the knots of h_n coincide with (some of) the order statistics $X_{(1)}, \dots, X_{(n)}$, introduce the set of knots $\hat{\mathcal{S}}_n(h_n)$ of h_n as

$$\hat{\mathcal{S}}_n(h_n) := \{t \in (X_{(1)}, X_{(n)}) : h'_n(t-) > h'_n(t+)\} \cup \{X_{(1)}, X_{(n)}\}.$$

Dümbgen and Rufibach (2009) found that $\hat{\varphi}_n$ is piecewise linear, that $\hat{\varphi}_n = -\infty$ on $\mathbb{R} \setminus [X_{(1)}, X_{(n)}]$ and that the knots of $\hat{\varphi}_n$ only occur at (some of the) ordered observations $X_{(1)} < \dots < X_{(n)}$. The latter property is entirely different from the estimation of a k -monotone density for $k > 1$ (see below), where the knots fall strictly between observations with probability equal to 1.

According to Theorem 2.4 in Dümbgen and Rufibach (2009), the estimator $\hat{\varphi}_n$ has the following characterization. For $x \geq X_{(1)}$ (recall that $\hat{\varphi}_n := -\infty$ outside $[X_{(1)}, X_{(n)}]$), define the processes

$$\begin{aligned} \hat{F}_n(x) &:= \int_{X_{(1)}}^x \exp(\hat{\varphi}_n(t)) dt, & \hat{H}_n(x) &:= \int_{X_{(1)}}^x \hat{F}_n(t) dt, \\ \mathbb{H}_n(x) &:= \int_{X_{(1)}}^x \mathbb{F}_n(t) dt = \int_{-\infty}^x \mathbb{F}_n(t) dt. \end{aligned}$$

Then, the concave function $\hat{\varphi}_n$ is the MLE of the log-density φ_0 if, and only if,

$$(1.1) \quad \hat{H}_n(x) \begin{cases} \leq \mathbb{H}_n(x), & \text{for all } x \geq X_{(1)}, \\ = \mathbb{H}_n(x), & \text{if } x \in \hat{\mathcal{S}}_n(\hat{\varphi}_n). \end{cases}$$

1.4. *Other shape constraints.* Maximum likelihood estimation of a monotone density f_0 on $[0, \infty)$ was first studied by Grenander (1956). Under the assumption that f_0 is C^1 in a neighborhood of a point $x_0 > 0$, such that $f'_0(x_0) < 0$,

Prakasa Rao (1969) established the (local) asymptotic distribution theory of the Grenander estimator \hat{f}_n :

$$n^{1/3}(\hat{f}_n(x_0) - f_0(x_0)) \xrightarrow{d} |f'_0(x_0)f_0(x_0)/2|^{1/3}\mathbb{Z},$$

where \mathbb{Z} is the slope at zero of the (least) concave majorant of the process $W(t) - t^2$, $t \in \mathbb{R}$ for two-sided Brownian motion W starting at 0.

Under the assumption that the true density f_0 is convex on $[0, \infty)$ and that f_0 is C^2 in a neighborhood of x_0 with $f''_0(x_0) > 0$, Groeneboom, Jongbloed and Wellner (2001b) show that the MLE \hat{f}_n (as well as the least squares estimator of f_0) satisfies

$$n^{2/5}(\hat{f}_n(x_0) - f_0(x_0)) \xrightarrow{d} (24^{-1}f_0^2(x_0)f''_0(x_0))^{1/5}\mathbb{H}''(0),$$

where \mathbb{H} is a particular upper envelope of an integrated two-sided Brownian motion $+t^4$ [see also Groeneboom, Jongbloed and Wellner (2001a)].

The classes of monotone and convex decreasing densities are particular cases of the class of k -monotone densities. Modulo a spline interpolation conjecture, Balabdaoui and Wellner (2007) were able to adapt the approach of Groeneboom, Jongbloed and Wellner (2001b) to this general class of densities.

We find that log-concave estimation shares many similarities with the aforementioned shape-constrained estimation problems. In particular, the limiting distribution of the MLE, our nonparametric estimator, involves a stochastic process whose second derivative is concave and which stays below an integrated Brownian motion minus t^{k+2} . The even integer k determines the number of vanishing derivatives of the true concave function φ_0 at the estimation point x_0 . Using Theorem 2.1, one can derive a procedure for estimation of k . This is relevant in practical applications of our results, that is, construction of confidence intervals for the mode using the limiting distribution given in Theorem 2.1. These problems are the subject of ongoing research.

1.5. Organization of the paper. In Section 2, we establish the limiting distributions of the ML estimators, $\hat{\varphi}_n$ and \hat{f}_n , at a fixed point $x_0 \in \mathbb{R}$ under some specified working assumptions. The characterization of either $\hat{\varphi}_n$ or \hat{f}_n given in (1.1) coincides, except for the direction of the inequality, with that of the least-squares estimator of a convex decreasing density, studied by Groeneboom, Jongbloed and Wellner (2001b); see their Lemma 2.2, page 1657. This enables us to adopt the general scheme of the proof in their paper.

Log-concave densities f and their logarithm φ can easily have vanishing second and higher derivatives at fixed points; an explicit example will be given in Section 2. Thus, the formulation of our asymptotic results allows higher derivatives of the concave function φ_0 to vanish at the estimation point. This is somewhat more general than the assumptions of Groeneboom, Jongbloed and Wellner (2001b) (where a natural assumption is that the second derivative is positive

at the point of interest, but similar vanishing of second derivatives and existence of a nonzero higher order derivative can also easily occur), but it is analogous to the results of Wright (1981) and Leurgans (1982) for nonparametric estimation of a monotone regression function. Similar results for the Grenander estimator of a monotone density are stated by Anevski and Hössjer (2006). We find that the respective limiting distributions of the MLE and its first derivative depend on a stochastic process, H_k , equal almost surely to the “lower envelope” (or just “invelope”) on \mathbb{R} of the integrated Brownian motion minus t^{k+2} , where k is the order of the first nonzero derivative of φ_0 at the point of interest.

In Section 3, the estimation point x_0 is taken to be equal to the mode, m_0 , defined to be the smallest point in the modal interval of the log-concave density f_0 . A natural estimator of m_0 , which we denote by \widehat{M}_n , can be taken to be the smallest number maximizing the MLE $\widehat{\varphi}_n$ or, equivalently, the smallest number maximizing the MLE \widehat{f}_n . In this section, we establish our second main result: the asymptotic distribution of \widehat{M}_n . Under the assumption that the second derivative $f_0''(m_0) < 0$, we show that this distribution depends on the random variable defined to be the argmax or mode of $H_2^{(2)}$ on \mathbb{R} . When the second, third and higher derivatives of order $k - 1$ or lower vanish at m_0 but $f_0^{(k)}(m_0) < 0$, then the limit distribution depends on the mode of $H_k^{(2)}$.

Proofs are deferred to Section 4.

To illustrate all the quantities for which we provide limiting distributions, in Figure 1 we give plots of \widehat{f}_n , $\widehat{\varphi}_n$, \widehat{F}_n and $\widehat{\lambda}_n = \widehat{f}_n / (1 - \widehat{F}_n)$, based on two samples of sizes $n = 20$ and $n = 200$ drawn from a Gamma(2, 1) density $f_0(x) = xe^{-x}1_{[0, \infty)}(x)$. All these plots were generated using the R-package logcondens [see Rufibach and Dümbgen (2007)].

2. Limiting distribution theory. To state the main result, we make the following assumptions.

2.1. *Assumptions.* Fix $x_0 \in \mathbb{R}$. We suppose that the true density $f_0 = \exp \varphi_0$ satisfies the following assumptions:

(A1) The density function $f_0 \in \mathcal{LC}$.

(A2) $f_0(x_0) > 0$.

(A3) The function φ_0 is at least twice continuously differentiable in a neighborhood of x_0 .

(A4) If $\varphi_0''(x_0) \neq 0$, then $k = 2$. Otherwise, suppose that k is the smallest integer such that $\varphi_0^{(j)}(x_0) = 0$, $j = 2, \dots, k - 1$, and $\varphi_0^{(k)}(x_0) \neq 0$, and $\varphi_0^{(k)}$ is continuous in a neighborhood of x_0 .

Note that concavity of φ_0 and (A3) and (A4) imply that k is necessarily even and that $\varphi_0^{(k)}(x_0) < 0$. Indeed, suppose that $k > 2$. Using Taylor expansion of φ_0''




FIG. 1. Examples for log-concave density, log-density, CDF, and hazard rate estimation for $n = 20, 200$ (--- true functions, — estimators). The dotted vertical lines indicate the set $\hat{\mathcal{S}}_n(\hat{\varphi}_n)$. The - - - vertical lines are placed at the mode of the estimated density.

up to degree $k - 2$, there exists a small $h > 0$ for which we can write

$$\varphi_0''(x) = \frac{\varphi_0^{(k)}(x_0)}{(k - 2)!} (x - x_0)^{k-2} + o((x - x_0)^{k-2}), \quad x \in [x_0 - h, x_0 + h].$$

Since $\varphi_0''(x) \leq 0$ for all $x \in [x_0 - h, x_0 + h]$, it follows that $k - 2$ is even [i.e., k is even and $\varphi_0^{(k)}(x_0) < 0$].

2.2. *Notation.* Let W denote two-sided Brownian motion, starting at 0. For $t \in \mathbb{R}$, define:

$$(2.1) \quad Y_k(t) = \begin{cases} \int_0^t W(s) ds - t^{k+2}, & \text{if } t \geq 0, \\ \int_t^0 W(s) ds - t^{k+2}, & \text{if } t < 0. \end{cases}$$

For the uniform norm of a bounded function f , we write $\|f\|_\infty = \sup_{x \in \mathbb{R}} |f(x)|$. The derivative of $\widehat{\varphi}_n$ at $x \in \mathbb{R}$ is as usual denoted by $\widehat{\varphi}'_n(x)$. However, if $x \in \widehat{\mathcal{J}}_n(\widehat{\varphi}_n)$, then we define $\widehat{\varphi}'_n(x)$ as the left-derivative.

THEOREM 2.1. *Suppose that (A1)–(A4) hold. Then,*

$$\left(\begin{matrix} n^{k/(2k+1)}(\widehat{f}_n(x_0) - f_0(x_0)) \\ n^{(k-1)/(2k+1)}(\widehat{f}'_n(x_0) - f'_0(x_0)) \end{matrix} \right) \xrightarrow{d} \left(\begin{matrix} c_k(x_0, \varphi_0) H_k^{(2)}(0) \\ d_k(x_0, \varphi_0) H_k^{(3)}(0) \end{matrix} \right)$$

and

$$\left(\begin{matrix} n^{k/(2k+1)}(\widehat{\varphi}_n(x_0) - \varphi_0(x_0)) \\ n^{(k-1)/(2k+1)}(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0)) \end{matrix} \right) \xrightarrow{d} \left(\begin{matrix} C_k(x_0, \varphi_0) H_k^{(2)}(0) \\ D_k(x_0, \varphi_0) H_k^{(3)}(0) \end{matrix} \right),$$

where H_k is the “lower envelope” of the process Y_k ; that is,

$H_k(t) \leq Y_k(t)$ for all $t \in \mathbb{R}$;

$H_k^{(2)}$ is concave;

$H_k(t) = Y_k(t)$, if the slope of $H_k^{(2)}$ decreases strictly at t .

The constants c_k, d_k, C_k and D_k are given by

$$(2.2) \quad c_k(x_0, \varphi_0) = \left(\frac{f_0(x_0)^{k+1} |\varphi_0^{(k)}(x_0)|}{(k + 2)!} \right)^{1/(2k+1)},$$

$$(2.3) \quad d_k(x_0, \varphi_0) = \left(\frac{f_0(x_0)^{k+2} |\varphi_0^{(k)}(x_0)|^3}{[(k + 2)!]^3} \right)^{1/(2k+1)},$$

$$(2.4) \quad C_k(x_0, \varphi_0) = \left(\frac{|\varphi_0^{(k)}(x_0)|}{f_0(x_0)^k (k + 2)!} \right)^{1/(2k+1)},$$

$$(2.5) \quad D_k(x_0, \varphi_0) = \left(\frac{|\varphi_0^{(k)}(x_0)|^3}{f_0(x_0)^{k-1} [(k + 2)!]^3} \right)^{1/(2k+1)}.$$

COROLLARY 2.2. *Suppose that (A1)–(A4) hold with $k = 2$. Then,*

$$\begin{pmatrix} n^{2/5}(\widehat{f}_n(x_0) - f_0(x_0)) \\ n^{1/5}(\widehat{f}'_n(x_0) - f'_0(x_0)) \end{pmatrix} \xrightarrow{d} \begin{pmatrix} c_2(x_0, \varphi_0)H_2^{(2)}(0) \\ d_2(x_0, \varphi_0)H_2^{(3)}(0) \end{pmatrix}$$

and

$$\begin{pmatrix} n^{2/5}(\widehat{\varphi}_n(x_0) - \varphi_0(x_0)) \\ n^{1/5}(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0)) \end{pmatrix} \xrightarrow{d} \begin{pmatrix} C_2(x_0, \varphi_0)H_2^{(2)}(0) \\ D_2(x_0, \varphi_0)H_2^{(3)}(0) \end{pmatrix},$$

where H_2 is the (concave) envelope of the process Y_2 ; that is,

$H_2(t) \leq Y_2(t)$ for all $t \in \mathbb{R}$;

$H_2^{(2)}$ is concave;

$H_2(t) = Y_2(t)$ if the slope of $H_2^{(2)}$ decreases strictly at t .

The constants c_2, d_2, C_2 and D_2 are given by (2.2)–(2.5), with $k = 2$.

Note that the constants $C_2(x_0, \varphi_0)$ and $D_2(x_0, \varphi_0)$, up to inversion of $f_0(x_0)$, exhibit a structure very similar to that of the constants given by Groeneboom, Jongbloed and Wellner (2001b) in the problem of estimating a convex density g_0 on $[0, \infty)$. We recall here that, in the latter problem, those constants are found to be equal to (we use our notation to make the comparison easy)

$$c_2(x_0, g_0) = \left(\frac{g_0(x_0)^2 g_0^{(2)}(x_0)}{4!} \right)^{1/5}, \quad d_2(x_0, g_0) = \left(\frac{g_0(x_0)(g_0^{(2)}(x_0))^3}{(4!)^3} \right)^{1/5}.$$

It is clear that φ_0 in the log-concave problem plays exactly the same role as f_0 in the problem of estimating a convex density. However, in the first case estimation is based on observations which are distributed according to $\exp \varphi_0$, whereas in the latter the data come from f_0 itself. A good insight into the difference between the expressions of the asymptotic constants can be gained from the proof of Theorem 4.6 in Section 4. There, we show that the leading coefficient of the drift of the limiting process Y_k depends on $\varphi_0^{(k)}(x_0) f_0(x_0) = f_0^{(k)}(x_0) - (\varphi_0'(x_0))^k f_0(x_0)$, where the second term is “filtered out” in the Taylor expansion of the estimation error in the neighborhood of x_0 . Hence, $|\varphi_0^{(k)}(x_0)| \cdot f_0(x_0)$ can be viewed as the dominating term replacing $|g_0^{(k)}(x_0)|$ in the convex estimation problem. For $k = 2$, the constants $c_2(x_0, \varphi_0)$ and $d_2(x_0, \varphi_0)$ given in (2.2) and (2.3), with $k = 2$, match closely with $c_2(x_0, g_0)$ and $d_2(x_0, g_0)$ obtained by Groeneboom, Jongbloed and Wellner (2001b) in the convex estimation problem, with $f_0(x_0)$ in the numerator, whereas $f_0(x_0)$ shows up in the denominator in the asymptotic constants $C_2(x_0, \varphi_0)$ and $D_2(x_0, \varphi_0)$. This results from applying the delta-method to $\widehat{f}_n(x_0) = \exp(\widehat{\varphi}_n(x_0))$ and $\widehat{f}'_n(x_0) = \widehat{\varphi}'_n(x_0) \widehat{f}_n(x_0)$, which yields $C_2(x_0, \varphi_0)$ and $D_2(x_0, \varphi_0)$.

Here is an explicit example showing how vanishing second (and higher) derivatives can occur. Consider the density function

$$f_0(x) = \sqrt{2} \frac{\Gamma(3/4)}{\pi} \exp(-x^4), \quad x \in \mathbb{R}.$$

In this case $\varphi_0^{(j)}(x_0) = 0, j = 1, 2, 3$ for $x_0 = 0$, and $\varphi_0^{(4)}(x_0) \neq 0$. The following “tilted” version of f_0 shows that vanishing second derivatives of φ_0 can also occur at points other than the mode of f :

$$\tilde{f}_0(x) = \exp(a + bx) f_0(x) = \tilde{a} \exp(bx - x^4),$$

where $\tilde{a} = \tilde{a}(b) := 1/\int_{\mathbb{R}} \exp(bx - x^4) dx$; in this case, $\tilde{\varphi}_0 := \log \tilde{f}_0$ satisfies $\tilde{\varphi}_0''(0) = 0$, but the mode $\tilde{m}_0 := M(\tilde{f}_0) = (b/4)^{1/3} > 0$ when $b > 0$, and $\tilde{\varphi}_0''(\tilde{m}_0) = -12(b/4)^{2/3} < 0$.

Finally, and in order to compare also the random parts of the limits in the convex and log-concave estimation problems, we would like to note that for our lower envelope process $H_k, -H_k$ has the same distribution as the “upper envelope” of $-Y_k$, which was called just the “invelope” in the case $k = 2$ by Groeneboom, Jongbloed and Wellner (2001b): The process $-Y_k$ has a drift equal to plus t^{k+2} , which specializes to t^4 in the convex density problem with $k = 2$. This “upper envelope” stays above $-Y_k$ and admits a convex second derivative. Since $-W$ has the same distribution as W , it follows that the upper and lower envelopes \mathbb{H}_k and H_k (associated with estimation of convex and concave functions, resp.) satisfy $\mathbb{H}_k \stackrel{d}{=} -H_k$. Since the derivatives at zero $\mathbb{H}_k^{(2)}(0)$ and $\mathbb{H}_k^{(3)}(0)$ of \mathbb{H}_k are distributed symmetrically about zero, the same is true of the derivatives at zero $H_k^{(2)}(0)$ and $H_k^{(3)}(0)$ of H_k .

As shown by Barlow and Proschan (1975), Lemma 5.8, page 77 [see also Marshall and Olkin (1979), page 493; Marshall and Olkin (2007), page 102; An (1998) and Bagnoli and Bergstrom (2005)], if f_0 is log-concave, then the hazard function

$$\lambda_0(x) = \frac{f_0(x)}{1 - F_0(x)} \mathbf{1}_{\{x < F_0^{-1}(1)\}}$$

is monotone nondecreasing. Defining the estimator of λ_0 based on \hat{f}_n as

$$\hat{\lambda}_n(x) = \frac{\hat{f}_n(x)}{1 - \hat{F}_n(x)} \mathbf{1}_{\{x < X_{(n)}\}},$$

application of the delta-method yields the following corollary.

COROLLARY 2.3. *Suppose that (A1)–(A4) hold. Then,*

$$\begin{pmatrix} n^{k/(2k+1)} (\hat{\lambda}_n(x_0) - \lambda_0(x_0)) \\ n^{(k-1)/(2k+1)} (\hat{\lambda}'_n(x_0) - \lambda'_0(x_0)) \end{pmatrix} \xrightarrow{d} \begin{pmatrix} g_k(x_0, \varphi_0) H_k^{(2)}(0) \\ h_k(x_0, \varphi_0) H_k^{(3)}(0) \end{pmatrix},$$

where the constants g_k and h_k are given by

$$g_k(x_0, \varphi_0) = c_k(x_0, \varphi_0)/(1 - F_0(x_0))$$

$$h_k(x_0, \varphi_0) = d_k(x_0, \varphi_0)/(1 - F_0(x_0)).$$

3. Inference about the mode of f_0 . Estimation of the mode of a unimodal density has been considered by many authors [see, e.g., Parzen (1962), Chernoff (1964), Grenander (1965), Dalenius (1965), Venter (1967), Wegman (1970a, 1970b, 1971), Eddy (1980, 1982), Hall (1982), Müller (1989), Romano (1988), Vieu (1996) and, more recently, Meyer (2001) and Herrmann and Ziegler (2004)].

Empirical studies of the performance of various estimators are given by Dalenius (1965), Ekblom (1972), Meyer (2001) and Meyer and Woodroffe (2004). Many of the methods considered for estimating the mode of a unimodal smooth density use kernel estimation, but others are based on the principle of substitution with another choice of estimator of the population density. For example, the estimators of Venter (1967) are related to nearest-neighbor estimators of the density f_0 . All the estimators of the mode in the class of unimodal densities known to us involve some more or less ad hoc choice, essentially because the maximum likelihood estimator of a unimodal density is not well defined, as explained by Birgé (1997). [Note that Wegman (1970b, 1971) discussed the nonparametric MLE of a unimodal density subject to a constraint on the height of the mode; without some constraint of this type, the MLE does not exist.]

For virtually all of the estimators of which we are aware, some choice of a smoothing parameter, bandwidth or constraint is required. Empirical choice of smoothing parameters has been studied by Müller (1989), who studied local methods of choosing the smoothing parameter, Grund and Hall (1995), who studied bootstrap methods, and Ziegler (2004), who studied plug-in methods. Klemelä (2005) gave a construction of adaptive estimators based on Lepski's method [Lepskiĭ (1991, 1992)]. For nonparametric Bayes estimators of unimodal densities and, hence, of the mode [see Brunner and Lo (1989) and Ho (2006a, 2006b)]; for these estimators, choice of a prior is equivalent to a choice of smoothing parameters.

In contrast, estimation in the (large) subclass of log-concave (or strongly unimodal) densities is much simpler, avoiding bandwidth or smoothing parameter choices completely. Since the maximum likelihood estimator exists, we can simply estimate the mode by the mode (or smallest point in a modal interval) of the MLE \hat{f}_n . Using the notation introduced by Eddy (1982) [and also used by Romano (1988)], we let $\tilde{M}_n := M(\hat{f}_n)$ where M denotes the mode functional (or “smallest argmax” functional) given by

$$M(g) := \min \left\{ t : g(t) = \max_{u \in \mathbb{R}} g(u) \right\}.$$

Because of the adaptive properties of the MLE’s \widehat{f}_n of f_0 and $\widehat{\varphi}_n$ of φ_0 , discussed in Section 1, we expect \widehat{M}_n to adapt to different local smoothness (or peakedness) hypotheses on f_0 [much as the Grenander estimator is locally adaptive in the case of estimating a monotone density, see, for example, Birgé (1989), page 1535]. Here, we study \widehat{M}_n as an estimator of the mode $M(f_0) := m_0$ under just the condition that f_0 has a continuous second derivative f_0'' in a neighborhood of m_0 , with $f_0''(m_0) < 0$. We begin in the next subsection with a new asymptotic minimax lower bound for estimation of m_0 under this hypothesis. The following subsection gives our new limiting distribution result for the MLE \widehat{M}_n of the mode m_0 .

3.1. *New lower bounds for estimating the mode.* Has’minskiĭ (1979) established a lower bound for estimation of the mode m_0 of a unimodal density $f \in \mathcal{U}$, assuming that f satisfies $f''(m_0) < 0$. He showed that the best local asymptotic minimax rate of convergence for any estimator of m_0 is $n^{-1/5}$. Has’minskiĭ based his proof on a sequence of parametric submodels of the form

$$f_n(x, \theta) = f(x) + \theta n^{-2/5} g(n^{1/5}(x - m_0)),$$

where, for $a := -f''(m_0)$,

$$g(x) := g_a(x) = \begin{cases} x, & \text{if } |x| \leq 1/a, \\ 0, & \text{if } |x| \geq K > 1/a \end{cases}$$

and $g := g_a$ satisfies $g(-x) = -g(x)$ and $|g''(x)| < a/2$ for all $x \in \mathbb{R}$. However, Has’minskiĭ (1979) did not study the dependence of the local minimax bound on $a = -f''(m_0)$ and $f(m_0)$, leaving his bound in terms of $c_0^2 := f(m_0) / \int g_a^2(x) dx$ involving the still unspecified function $g = g_a$.

Here, we consider different parametric submodels and derive the dependence of the constant in local asymptotic minimax lower bound for estimation of the mode m_0 in the family \mathcal{LC} of log-concave (or strongly unimodal) densities.

We want to derive asymptotic lower bounds for the local minimax risks for estimating the mode $M(f)$. The L_1 -minimax risk for estimating a functional ν of f_0 , based on a sample X_1, \dots, X_n of size n from f_0 , which is known to be in a subset $\mathcal{LC}_{n,\tau}$ of \mathcal{LC} is defined by

$$(3.1) \quad MMR_1(n, T_n, \mathcal{LC}_{n,\tau}) := \inf_{T_n} \sup_{f \in \mathcal{LC}_{n,\tau}} E_f |T_n - \nu(f)|,$$

where the infimum ranges over all possible measurable functions $T_n = t_n(X_1, \dots, X_n)$ mapping \mathbb{R}^n to \mathbb{R} . The shrinking classes $\mathcal{LC}_{n,\tau}$ used here are Hellinger balls centered at f_0 :

$$\mathcal{LC}_{n,\tau} = \left\{ f \in \mathcal{LC} : H^2(f, f_0) = \frac{1}{2} \int_{-\infty}^{\infty} (\sqrt{f(z)} - \sqrt{f_0(z)})^2 dz \leq \tau/n \right\}.$$

Consider estimation of

$$(3.2) \quad \nu(f) := M(f) = \inf \left\{ t \in \mathbb{R} : t = \sup_{u \in \mathbb{R}} f(u) \right\}.$$

Let $f_0 \in \mathcal{LC}$ and $m_0 = M(f_0)$ be fixed, such that f_0 is twice continuously differentiable at m_0 and $f_0''(m_0) < 0$. Consider the family $\{\varphi_\varepsilon\}_{\varepsilon>0}$ and resulting family $\{f_\varepsilon\}_{\varepsilon>0}$, defined as follows much as:

$$\varphi_\varepsilon(x) = \begin{cases} \varphi_0(x), & x < m_0 - \varepsilon c_\varepsilon, \\ \varphi_0(x), & x > m_0 + \varepsilon, \\ \varphi_0(m_0 + \varepsilon), & \\ \quad + \varphi_0'(m_0 + \varepsilon)(x - m_0 - \varepsilon), & x \in [m_0 - \varepsilon, m_0 + \varepsilon], \\ \varphi_0(m_0 - \varepsilon c_\varepsilon), & \\ \quad + \varphi_0'(m_0 - \varepsilon c_\varepsilon)(x - m_0 + \varepsilon c_\varepsilon), & x \in [m_0 - \varepsilon c_\varepsilon, m_0 - \varepsilon], \end{cases}$$

where c_ε is chosen so that φ_ε is continuous at $m_0 - \varepsilon$. Note that if $\varphi_0(x) = \gamma - \gamma_0(x - m_0)^2$, then $c_\varepsilon = 3$, for all ε , and $c_\varepsilon \rightarrow 3$, as $\varepsilon \downarrow 0$, since $f_0''(m_0) < 0$. Now define

$$h_\varepsilon(x) := \exp(\varphi_\varepsilon(x)) \quad \text{and} \quad f_\varepsilon(x) := \frac{h_\varepsilon(x)}{\int h_\varepsilon(y) dy}.$$

Then, f_ε is log-concave for each $\varepsilon > 0$ with mode $m_0 - \varepsilon$ by construction, so with $\nu(f_\varepsilon) := M(f_\varepsilon) :=$ the mode of f_ε , we have

$$\nu(f_\varepsilon) - \nu(f_0) = M(f_\varepsilon) - M(f_0) = m_0 - \varepsilon - m_0 = -\varepsilon.$$

Furthermore, the following lemma holds.

LEMMA 3.1. *Under the above assumptions,*

$$H^2(f_\varepsilon, f_0) = \frac{2f_0''(m_0)^2}{5f_0(m_0)}\varepsilon^5 + o(\varepsilon^5) := \rho\varepsilon^5 + o(\varepsilon^5).$$

PROOF. Proceeding as in [Jongbloed \(1995\)](#),

$$\begin{aligned} H^2(f_\varepsilon, f_0) &= \frac{1}{2} \int_{-\infty}^{\infty} [\sqrt{f_\varepsilon(x)} - \sqrt{f_0(x)}]^2 dx \\ &= \frac{1}{2} \int_{m_0 - \varepsilon c_\varepsilon}^{m_0 + \varepsilon} [\sqrt{f_\varepsilon(x)} - \sqrt{f_0(x)}]^2 dx \\ &= \frac{2}{5} f_0(m_0) \varphi_0''(m_0)^2 \varepsilon^5 + o(\varepsilon^5) = \frac{2}{5} \frac{f_0''(m_0)^2}{f_0(m_0)} \varepsilon^5 + o(\varepsilon^5) \end{aligned}$$

as $\varepsilon \downarrow 0$. Calculations similar to those of [Jongbloed \(1995\)](#) [see also [Jongbloed \(2000\)](#) and [Groeneboom, Jongbloed and Wellner \(2001b\)](#)] complete the proof of the lemma. \square

Taking $\varepsilon = cn^{-1/5}$ and defining $f_n := f_{cn^{-1/5}}$ yields

$$v(f_n) - v(f_0) = M(f_n) - M(f_0) = -cn^{-1/5}$$

and

$$nH^2(f_n, f_0) = \frac{2}{5} \frac{f_0''(m_0)^2}{f_0(m_0)} c^5 + o(1) := \rho c^5 + o(1).$$

Plugging these into the lower bound Lemma 4.1 of Groeneboom (1996), with $\ell(x) := |x|$, yields

$$\begin{aligned} & \liminf_n \inf_{T_n} n^{1/5} \max\{E_{n, P_n} |T_n - M(f_n)|, E_{n, P} |T_n - M(f_0)|\} \\ & \geq \frac{1}{4} c \exp(-2\rho c^5) = \frac{e^{-1/5}}{4 \cdot 10^{1/5}} \rho^{-1/5} = (0.15512) \left(\frac{f_0(m_0)}{f_0''(m_0)^2} \right)^{1/5} \end{aligned}$$

by choosing $c = (10\rho)^{-1/5}$. This yields the following proposition.

PROPOSITION 3.2 (Minimax risk lower bound). *Suppose that $v(f) = M(f)$, as defined in (3.2), and that $\mathcal{LC}_{n,\tau}$ is as defined above where f_0'' is continuous in a neighborhood of $m_0 = M(f_0)$ with $f_0''(m_0) < 0$. Then,*

$$\begin{aligned} & \sup_{\tau > 0} \limsup_{n \rightarrow \infty} n^{1/5} \inf_{T_n} \sup_{f \in \mathcal{LC}_{n,\tau}} E_f |T_n - M(f)| \\ & \geq \left(\frac{5/2}{4^5 \cdot e \cdot 10} \right)^{1/5} \left(\frac{f_0(m_0)}{f_0''(m_0)^2} \right)^{1/5} \doteq (0.15512) \left(\frac{f_0(m_0)}{f_0''(m_0)^2} \right)^{1/5}. \end{aligned}$$

REMARK 3.3. Note that the constant $b(f_0, m_0) := (f_0(m_0)/f_0''(m_0)^2)^{1/5}$ appearing on the right-hand side of this lower bound is scale equivariant in exactly the right way: if $f_c(x) := f_0(m_0 + (x - m_0)/c)/c$ for $c > 0$, then $b(f_c, m_0) = cb(f_0, m_0)$ for all $c > 0$. The constant $b(f_0, m_0)$ will appear in the limit distribution appearing in the next subsection.

REMARK 3.4. If \mathcal{LC} is replaced by the class \mathcal{U} of unimodal densities on \mathbb{R} and $\mathcal{LC}_{n,\tau}$ is replaced by $\mathcal{U}_{n,\tau}$ defined analogously where f_0 satisfies $f_0''(m_0) < 0$ and f_0'' continuous in a neighborhood of m_0 , then a minimax lower bound of the same form as Proposition 3.2 holds with exactly the same dependence on $b(f_0, m_0) = (f_0(m_0)/f_0''(m_0)^2)^{1/5}$, but with the absolute constant $0.15512\dots$ replaced by $0.19784\dots$. This can be seen by taking the perturbations $\{f_\varepsilon\}_{\varepsilon > 0}$ defined by

$$f_\varepsilon(x) = \begin{cases} f_0(x), & x \leq x_0 - \varepsilon, \\ f_0(x), & x > x_0 + \varepsilon, \\ f_0(x_0) + b_\varepsilon(x - x_0 + \varepsilon), & x_0 - \varepsilon \leq x \leq x_0 + \varepsilon, \end{cases}$$

where b_ε is chosen so that $f_\varepsilon(x_0 + \varepsilon) > f_0(x_0 + \varepsilon)$ and $\int_{x_0-\varepsilon}^{x_0+\varepsilon} f_\varepsilon(x) dx = \int_{x_0-\varepsilon}^{x_0+\varepsilon} f_0(x) dx$.

REMARK 3.5. If φ_0 is continuously k -times differentiable in a neighborhood of the mode m_0 , $\varphi_0^{(j)}(m_0) = 0$ for $j = 2, \dots, k - 1$ and $\varphi_0^{(k)}(m_0) \neq 0$ [assumption (A4)], then it can be shown that the minimax rate of convergence is $n^{1/(2k+1)}$ and that the minimax lower bound is proportional to

$$\left(\frac{1}{f_0(m_0)\varphi_0^{(k)}(m_0)^2} \right)^{1/(2k+1)} = \left(\frac{f_0(m_0)}{f_0^{(k)}(m_0)^2} \right)^{1/(2k+1)},$$

where the proportionality constant depends on the largest root of the polynomial $x^k - (k/(k - 1))x^{k-1} - (2k - 1)/(k - 1)$ (which equals 3 when $k = 2$).

3.2. *Limiting distribution for the MLE \widehat{M}_n in \mathcal{LC} .* Now, let \widehat{f}_n be the MLE of f in the class \mathcal{LC} of log-concave densities, and let $\widehat{M}_n = M(\widehat{f}_n)$, $m_0 = M(f_0)$. Here is our result concerning the limiting distribution of \widehat{M}_n under the same assumptions on f_0 as in the previous section on lower bounds.

THEOREM 3.6. *Suppose that f_0'' is continuous in a neighborhood of $m_0 = M(f_0)$ and that $f_0''(m_0) < 0$. Then,*

$$n^{1/5}(\widehat{M}_n - m_0) \xrightarrow{d} \left(\frac{(4!)^2 f_0(m_0)}{f_0''(m_0)^2} \right)^{1/5} M(H_2^{(2)}).$$

Note that the limiting distribution depends on a multiple of the same constant $b(f_0, m_0)$, which appears in the asymptotic minimax lower bound of Proposition 3.2, times a universal term $M(H_2^{(2)})$, the mode of the “estimator” $H_2^{(2)}(t)$ of the canonical concave function $-12t^2$ in the limit Gaussian problem: estimate the mode of $f_0(t) = -12t^2$, based on observation of $Y(t) = \int_0^t X(s) ds$, when

$$dX(t) = f_0(t) dt + dW(t).$$

We expect that this distribution, namely the distribution of

$$M(H_2^{(2)}) = \arg \max_{t \in \mathbb{R}} H_2^{(2)}(t),$$

will occur in several other problems involving nonparametric estimation of the mode or antimode of convex or concave functions under similar second derivative hypotheses. For example, it seems clear that it will occur as the limiting distribution of the nonparametric estimator of the antimode of a convex bathtub-shaped hazard [in the setting of Jankowski and Wellner (2007)]; as the limiting distribution of the nonparametric estimator of the antimode of a convex regression function in the setting of Groeneboom, Jongbloed and Wellner (2001b); and as the limiting distribution of the nonparametric estimator of the mode of a concave regression function.

When $\varphi_0^{(j)}(m_0) = 0$, for $j = 2, \dots, k - 1$, $\varphi_0^{(k)}(m_0) \neq 0$, and $\varphi_0^{(k)}$ is continuous in a neighborhood of m_0 , then an analogous result (with a completely similar proof) holds:

$$n^{1/(2k+1)}(\widehat{M}_n - m_0) \xrightarrow{d} \left(\frac{(k+2)!^2}{f_0(m_0)|\varphi_0^{(k)}(m_0)|^2} \right)^{1/(2k+1)} M(H_k^{(2)}).$$

In particular, when $k = 4$, the rate of convergence is $n^{1/9}$, and the limit distribution becomes that of

$$\left(\frac{6!^2 f_0(m_0)}{f_0^{(4)}(m_0)^2} \right)^{1/9} M(H_4^{(2)}).$$

Apparently, estimation of m_0 becomes considerably more difficult when the second and possibly higher order derivatives of φ_0 vanish at m_0 .

On the other hand, if φ_0 (or equivalently, f_0) is cusp-shaped at m_0 , then the rate of convergence of \widehat{M}_n is $n^{1/3}$, and the local asymptotic minimax rate of convergence is also $n^{1/3}$; we will pursue these issues elsewhere.

4. Proofs for Sections 2 and 3. Throughout this section, we fix k and let

$$\begin{aligned} r_n &:= n^{(k+2)/(2k+1)}, & s_n &:= n^{-1/(2k+1)}, \\ x_n(t) &:= x_{n,k}(t) := x_0 + s_n t := x_0 + n^{-1/(2k+1)}t, \\ \mathfrak{I} &:= \mathfrak{I}(x_0, n, k, t) := \begin{cases} [x_0, x_n(t)], & t \geq 0, \\ [x_n(t), x_0], & t < 0. \end{cases} \end{aligned}$$

4.1. *Preparation: technical lemmas and tightness results.* First, some notation.

Local processes: The local processes $\mathbb{Y}_n^{\text{loc}}$ and $\widehat{H}_n^{\text{loc}}$ are defined for $t \in \mathbb{R}$ by

$$\mathbb{Y}_n^{\text{loc}}(t) := r_n \int_{x_0}^{x_n(t)} \left(\mathbb{F}_n(v) - \mathbb{F}_n(x_0) - \int_{x_0}^v \left(\sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) du \right) dv$$

and

$$\begin{aligned} \widehat{H}_n^{\text{loc}}(t) &:= r_n \int_{x_0}^{x_n(t)} \int_{x_0}^v \left(\widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) du dv \\ &\quad + \widehat{A}_n t + \widehat{B}_n, \end{aligned}$$

where in the limit Gaussian problem: estimate the mode

$$(4.1) \quad \widehat{A}_n = r_n s_n (\widehat{F}_n(x_0) - \mathbb{F}_n(x_0)) \quad \text{and}$$

$$(4.2) \quad \widehat{B}_n = r_n (\widehat{H}_n(x_0) - \mathbb{H}_n(x_0)).$$

We also define the “modified” local processes

$$\begin{aligned}
 \mathbb{Y}_n^{\text{locmod}}(t) := & \frac{r_n}{f_0(x_0)} \int_{x_0}^{x_n(t)} \left(\mathbb{F}_n(v) - \mathbb{F}_n(x_0) \right. \\
 (4.3) \quad & \left. - \int_{x_0}^v \left(\sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) du \right) dv \\
 & - r_n \int_{x_0}^{x_n(t)} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv
 \end{aligned}$$

and

$$\begin{aligned}
 \mathbb{H}_n^{\text{locmod}}(t) := & r_n \int_{x_0}^{x_n(t)} \int_{x_0}^v (\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi_0'(x_0)) du dv \\
 (4.4) \quad & + \frac{\widehat{A}_n t + \widehat{B}_n}{f_0(x_0)},
 \end{aligned}$$

where $\widehat{\Psi}_{k,n,2}$ is defined below in (4.26).

The following lemma uses the notion of uniform covering numbers [see [van der Vaart and Wellner \(1996\)](#), Sections 2.1 and 2.7] for complete definitions and further information.

LEMMA 4.1. *Let \mathcal{F} be a collection of functions defined on $[x_0 - \delta, x_0 + \delta]$, with $\delta > 0$ small and let $s > 0$. Suppose that for a fixed $x \in [x_0 - \delta, x_0 + \delta]$ and $R > 0$, such that $[x, x + R] \subseteq [x_0 - \delta, x_0 + \delta]$, the collection*

$$\mathcal{F}_{x,R} = \{ f_{x,y} := f 1_{[x,y]}, f \in \mathcal{F}, x \leq y \leq x + R \}$$

admits an envelope $F_{x,R}$, such that

$$EF_{x,R}^2(X_1) \leq KR^{2d-1}, \quad R \leq R_0$$

for some $d \geq 1/2$ and $K > 0$, depending only on x_0 and δ . Moreover, suppose that

$$(4.5) \quad \sup_Q \int_0^1 \sqrt{\log N(\eta \|F_{x,R}\|_{Q,2}, \mathcal{F}_{x,R}, L_2(Q))} d\eta < \infty.$$

Then, for each $\varepsilon > 0$, there exist random variables M_n of order $O_p(1)$ (not depending on x or y) and $R_0 > 0$, such that

$$\left| \int f_{x,y} d(\mathbb{F}_n - F_0) \right| \leq \varepsilon |y - x|^{s+d} + n^{-(s+d)/(2s+1)} M_n \quad \text{for } |y - x| \leq R_0.$$

PROOF. See [Kim and Pollard \(1990\)](#) and [Balabdaoui and Wellner \(2007\)](#), Lemmas 4.4 and 6.1. The special case $s = 1 = d$ is Lemma 4.1 of [Kim and Pollard \(1990\)](#). \square

LEMMA 4.2. *If (A3) and (A4) hold, then*

$$(4.6) \quad f_0^{(j)}(x_0) = [\varphi_0'(x_0)]^j f_0(x_0) \quad \text{for } j = 1, \dots, k - 1$$

and, for $j = k$

$$f_0^{(k)}(x_0) = (\varphi_0^{(k)}(x_0) + [\varphi_0'(x_0)]^k) f_0(x_0).$$

PROOF. The expressions for $f_0^{(j)}(x_0)$ follow immediately from a recursive argument using the identity $f_0 = \exp \varphi_0$ and the assumption $\varphi_0^{(j)}(x_0) = 0$, for $j = 2, \dots, k - 1$, if $k > 2$. \square

Now, let $\tau_n^+ := \inf\{t \in \widehat{\mathcal{J}}(\widehat{\varphi}_n) : t > x_0\}$ and $\tau_n^- := \sup\{t \in \widehat{\mathcal{J}}(\widehat{\varphi}_n) : t < x_0\}$.

THEOREM 4.3. *If (A1)–(A4) hold, then*

$$(4.7) \quad \tau_n^+ - \tau_n^- = O_p(n^{-1/(2k+1)}).$$

Theorem 4.3 should be compared to Theorem 3.3 of Dümbgen and Rufibach (2009). When their Theorem 3.3 is specialized to the case $\beta = 2$, so that $\varphi_0''(x) \leq C < 0$, for all $x \in T := [A, B]$, then it yields the following: If m_n denotes the number of elements in $\mathcal{J}_n(\widehat{\varphi}_n) \cap T$, then for any successive knot points t_{i-1} and t_i in $\mathcal{J}_n(\widehat{\varphi}_n) \cap T$,

$$(4.8) \quad \sup_{i=2, \dots, m_n} (t_i - t_{i-1}) = O_p(\rho_n^{1/5}),$$

where $\rho_n = \log(n)/n$.

PROOF OF THEOREM 4.3. From the first characterization of the estimator \widehat{f}_n in Dümbgen and Rufibach (2009), for every function Δ such that $\widehat{\varphi}_n + t \Delta$ is concave for a $t > 0$ small enough, we know that

$$(4.9) \quad \int_{\mathbb{R}} \Delta(x) d\mathbb{F}_n(x) \leq \int_{\mathbb{R}} \Delta(x) d\widehat{F}_n(x).$$

This is equivalent to

$$(4.10) \quad \int_{\mathbb{R}} \Delta(x) d(\mathbb{F}_n(x) - F_0(x)) \leq \int_{\mathbb{R}} \Delta(x) (\widehat{f}_n(x) - f_0(x)) dx.$$

Using specific indicator functions for Δ , one can furthermore show that

$$(4.11) \quad \widehat{F}_n(\tau) \in [\mathbb{F}_n(\tau) - 1/n, \mathbb{F}_n(\tau)]$$

for every $\tau \in \widehat{\mathcal{J}}_n(\widehat{\varphi}_n)$ [see Rufibach (2006) and Corollary 2.5 of Dümbgen and Rufibach (2009)].

Now, the idea is to choose a particular permissible perturbation function Δ that satisfies the following two conditions:

1. Δ is “local,” that is, compactly supported on $[\tau_n^-, \tau_n^+]$.
2. Δ should “filter” out the unknown error $\widehat{f}_n - f_0$.

The second requirement means that Δ should be chosen so that

$$(4.12) \quad \int_{\tau_n^-}^{\tau_n^+} \Delta(x) dx = 0, \quad \int_{\tau_n^-}^{\tau_n^+} \Delta(x)(x - \bar{\tau}) dx = 0,$$

where $\bar{\tau} := (\tau_n^- + \tau_n^+)/2$ is the mid-point of $[\tau_n^-, \tau_n^+]$. If this is guaranteed, then the right-hand side of (4.10) in the end will only depend on the distance $\tau_n^+ - \tau_n^-$ and $f_0(x_0)$.

Define Δ_0 by

$$\Delta_0(x) = (x - \tau_n^-)1_{[\tau_n^-, \bar{\tau}]}(x) + (\tau_n^+ - x)1_{[\bar{\tau}, \tau_n^+]}(x).$$

Since $\widehat{\varphi}_n + t\Delta_0$ is concave for small $t > 0$, Δ_0 is permissible. It is also compactly supported. However, since Δ_0 is nonnegative, there is no hope that it fulfills the second of the requirements above. We therefore introduce a modified perturbation function

$$\Delta_1(x) = \Delta_0(x) - \frac{1}{4}(\tau_n^+ - \tau_n^-)1_{[\tau_n^-, \tau_n^+]}(x), \quad x \in \mathbb{R}.$$

Clearly, existence of a $t > 0$, such that $\widehat{\varphi}_n + t\Delta_1$ is concave, is no longer guaranteed. However, using (4.11),

$$\begin{aligned} & \int \Delta_1(x) d(\mathbb{F}_n - F_0)(x) \\ &= \int \Delta_1(x) d(\mathbb{F}_n - \widehat{F}_n)(x) + \int \Delta_1(x) d(\widehat{F}_n - F_0)(x) \\ (4.13) \quad & \leq \frac{\tau_n^+ - \tau_n^-}{4} \left| \int_{\tau_n^-}^{\tau_n^+} d(\mathbb{F}_n - \widehat{F}_n)(x) \right| + \int \Delta_1(x) d(\widehat{F}_n - F_0)(x) \end{aligned}$$

$$(4.14) \quad \leq \frac{\tau_n^+ - \tau_n^-}{2n} + \int \Delta_1(x)(\widehat{f}_n - f_0)(x) dx.$$

To get the inequality in (4.13), we used (4.9) with $\Delta = \Delta_0$ and (4.11). The next step is to get bounds for the integrals in the crucial inequality (4.14). Define

$$R_{1n} := \int \Delta_1(x)(\widehat{f}_n - f_0)(x) dx$$

and

$$R_{2n} := \int \Delta_1(x) d(\mathbb{F}_n - F_0)(x).$$

Rearranging the inequality in (4.14) and using these definitions yields

$$-R_{1n} \leq \frac{\tau_n^+ - \tau_n^-}{2n} - R_{2n}.$$

Consistency of $\widehat{\varphi}_n$, together with $\varphi_0^{(k)}(x_0) < 0$, implies $\tau_n^+ - \tau_n^- = o_p(1)$. Thus, it follows from Lemma 4.4 that

$$M_k(-\varphi_0^{(k)}(x_0))(\tau_n^+ - \tau_n^-)^{k+2}(1 + o_p(1)) \leq o_p(1)n^{-1} + O_p(r_n^{-1}) = O_p(r_n^{-1}).$$

This yields the claimed rate, $O_p(n^{-1/(2k+1)})$, for the distance between τ_n^+ and τ_n^- . \square

LEMMA 4.4. *Suppose (A1)–(A4) hold. Then,*

$$R_{2n} = O_p(r_n^{-1})$$

and

$$R_{1n} = M_k f_0(x_0)\varphi_0^{(k)}(x_0)(\tau_n^+ - \tau_n^-)^{k+2} + o_p((\tau_n^+ - \tau_n^-)^{k+2}),$$

where $M_k > 0$ depends only on k and $\varphi_0^{(k)}(x_0) < 0$.

PROOF. Define the function $p_n(t) = \widehat{\varphi}_n(t) - \varphi_0(t)$ for any $t \in [\tau_n^-, \tau_n^+]$. Then, using Taylor expansion of $h \mapsto \exp(h)$ up to order k , we can find $\theta_{t,n} \in [\tau_n^-, \tau_n^+]$, such that

$$R_{1n} = \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) \left(\sum_{j=1}^{k-1} \frac{p_n(t)^j}{j!} + \frac{1}{k!} \exp(\theta_{t,n}) p_n(t)^k \right) dt := \sum_{j=1}^k \frac{S_{nj}}{j!},$$

where

$$S_{nj} := \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) p_n(t)^j dt \quad \text{for } 1 \leq j \leq k - 1$$

and

$$S_{nk} := \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) f_0(t) \exp(\theta_{t,n}) p_n(t)^k dt.$$

If we expand $f_0(t)$ around the mid-point $\bar{\tau}$ of $[\tau_n^-, \tau_n^+]$, we get, for $1 \leq j \leq k - 1$ and a $\eta_{n,t,j} \in [\tau_n^-, \tau_n^+]$,

$$\begin{aligned} S_{nj} &= \sum_{l=0}^{k-1} \frac{f_0^{(l)}(\bar{\tau})}{l!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) (t - \bar{\tau})^l p_n(t)^j dt \\ &\quad + \int_{\tau_n^-}^{\tau_n^+} \frac{f_0^{(k)}(\eta_{n,t,j})}{k!} \Delta_1(t) (t - \bar{\tau})^k p_n(t)^j dt \end{aligned}$$

and, for $j = k$

$$\begin{aligned} S_{nk} &= \sum_{l=0}^{k-1} \frac{f_0^{(l)}(\bar{\tau})}{l!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) \exp(\theta_{t,n}) (t - \bar{\tau})^l p_n(t)^k dt \\ &\quad + \int_{\tau_n^-}^{\tau_n^+} \frac{f_0^{(k)}(\eta_{n,t,k})}{k!} \Delta_1(t) \exp(\theta_{t,n}) (t - \bar{\tau})^k p_n(t)^k dt. \end{aligned}$$

It turns out that the dominating term in R_{1n} is the first term in the Taylor expansion of S_{n1} . All the other terms are of smaller order since both p_n and $(t - \bar{\tau})^l$, $l > 0$, are $o_p(1)$ uniformly in $t \in [\tau_n^-, \tau_n^+]$. We denote this dominating term by Q_{n1} . Since $\widehat{\varphi}_n$ is linear on $[\tau_n^-, \tau_n^+]$, we write $\widehat{\varphi}_n(t) = \widehat{\varphi}_n(\bar{\tau}) + (t - \bar{\tau})\widehat{\varphi}'_n(\bar{\tau})$. By Taylor expansion of p_n around $\bar{\tau}$, we get

$$\begin{aligned} \frac{Q_{1n}}{f_0(\bar{\tau})} &= \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) p_n(t) dt \\ &= p_n(\bar{\tau}) \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t) dt + p'_n(\bar{\tau}) \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t)(t - \bar{\tau}) dt \\ &\quad - \sum_{j=2}^k \frac{\varphi_0^{(j)}(\bar{\tau})}{j!} \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t)(t - \bar{\tau})^j dt - \int_{\tau_n^-}^{\tau_n^+} \varepsilon_n(t) \Delta_1(t)(t - \bar{\tau})^k dt, \end{aligned}$$

where the first two terms are zero, since (4.12) holds when $\Delta = \Delta_1$ and $\|\varepsilon_n\|_\infty \rightarrow_p 0$ as $\tau_n^+ - \tau_n^- \rightarrow_p 0$. Using the fact that

$$(4.15) \quad \int_{\tau_n^-}^{\tau_n^+} \Delta_1(t)(t - \bar{\tau})^j dt = \begin{cases} 0, & \text{for } j = 0 \text{ and } j \text{ odd,} \\ (\tau_n^+ - \tau_n^-)^{j+2} \left(\frac{-j}{2^{(j+2)}(j+1)(j+2)} \right), & \text{for } j \text{ even,} \end{cases}$$

we conclude that

$$Q_{1n} = \frac{k}{2^{(k+2)}k!(k+1)(k+2)} f_0(\bar{\tau})\varphi_0^{(k)}(\bar{\tau})((\tau_n^+ - \tau_n^-)^{k+2} + o_p(1))$$

and the claimed form of R_{1n} in the lemma follows.

For R_{2n} , we proceed along the lines of the proof of Lemma 4.1 in [Groeneboom, Jongbloed and Wellner \(2001b\)](#). This means we have to line up with the assumption of Theorem 2.14.1 in [van der Vaart and Wellner \(1996\)](#). Therefore, define a generalized version of R_{2n} :

$$R_{2n}^{x,y} = \int_x^y \Delta_1(z) d(\mathbb{F}_n - F_0)(z)$$

for $-\infty < x \leq y$. With this function, we have, for some $R > 0$,

$$\begin{aligned} &\sup_{y: 0 \leq y-x \leq R} |R_{2n}^{x,y}| \\ &= 2 \sup_{y: 0 \leq y-x \leq R} \left| \int_x^{(x+y)/2} (z - x - \frac{1}{4}(y-x)) d(\mathbb{F}_n - F_0)(z) \right| \\ &= 2 \sup_{y: 0 \leq y-x \leq R} \left| \int h_{x,y}(z) d(\mathbb{F}_n - F_0)(z) \right|, \end{aligned}$$

where

$$h_{x,y}(z) = (z - x - \frac{1}{4}(y - x))1_{[x,(x+y)/2]}(z) = h(z)1_{[x,(x+y)/2]}(z).$$

Then, the collection of functions

$$\mathcal{F}_{x,R} = \{h1_{[x,(x+y)/2]} : x \leq y \leq x + R\}$$

is a Vapnik–Chervonenkis subgraph class with envelope function

$$F_{x,R}(z) = ((z - x) + R/4)1_{[x,x+R]}(z).$$

Finally, Theorem 2.6.7 in van der Vaart and Wellner (1996) yields the entropy condition (4.5).

A log-concave density is always unimodal and the value at the mode is finite, and hence, $K := \|f_0\|_\infty$ is finite. Therefore,

$$\begin{aligned} EF_{x,R}^2(X_1) &= \int_x^{x+R} (z - x)^2 f_0(z) dz + \frac{R}{2} \int_x^{x+R} (z - x) f_0(z) dz + \frac{R^2}{16} \int_x^{x+R} f_0(z) dz \\ &\leq \left(\frac{K}{3}(z - x)^3 + \frac{RK}{4}(z - x)^2 + \frac{R^2K}{16}z \right) \Big|_{z=x}^{x+R} \\ &= \frac{31}{48}KR^3. \end{aligned}$$

It follows from Lemma 4.1, with $d = 2$ and $s = k$, that $R_{2n} = O_p(r_n^{-1})$. \square

4.2. Proofs for Section 2.

LEMMA 4.5. For any $M > 0$, we have

$$(4.16) \quad \sup_{|t| \leq M} |\widehat{\varphi}'_n(x_0 + s_n t) - \varphi'_0(x_0)| = O_p(s_n^{k-1}),$$

$$(4.17) \quad \sup_{|t| \leq M} |\widehat{\varphi}_n(x_0 + s_n t) - \varphi_0(x_0) - s_n t \varphi'_0(x_0)| = O_p(s_n^k).$$

Furthermore, if we define, for any $u \in \mathbb{R}$,

$$\widehat{e}_n(u) = \widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j - f_0(x_0) \frac{[\varphi'_0(x_0)]^k}{k!} (u - x_0)^k,$$

then

$$(4.18) \quad \begin{aligned} &\sup_{|t| \leq M} |\widehat{e}_n(x_0 + s_n t) - f_0(x_0)(\widehat{\varphi}_n(x_0 + s_n t) - \varphi_0(x_0) - s_n t \varphi'_0(x_0))| \\ &= o_p(s_n^k). \end{aligned}$$

PROOF. The proof of (4.16) and (4.17) is identical to that of Lemma 4.4 in Groeneboom, Jongbloed and Wellner (2001b) since the characterization of \widehat{f}_n given in (1.1) is (up to the direction of the inequality) equivalent to that of the least-squares estimator of a convex density.

Now, we prove (4.18). Using Taylor expansion of $h \mapsto \exp(h)$ up to order k around zero, we can write

$$\begin{aligned}
 (4.19) \quad \widehat{f}_n(u) - f_0(x_0) &= f_0(x_0)[\exp(\widehat{\varphi}_n(u) - \varphi_0(x_0)) - 1] \\
 &= f_0(x_0) \sum_{j=1}^k \frac{1}{j!} (\widehat{\varphi}_n(u) - \varphi_0(x_0))^j + f_0(x_0) \widehat{\Psi}_{k,n,1}(u),
 \end{aligned}$$

where

$$\widehat{\Psi}_{k,n,1}(u) = \sum_{j=k+1}^{\infty} \frac{1}{j!} (\widehat{\varphi}_n(u) - \varphi_0(x_0))^j.$$

But, for any $j \geq 1$,

$$\begin{aligned}
 (4.20) \quad &(\widehat{\varphi}_n(u) - \varphi_0(x_0))^j \\
 &= [\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi'_0(x_0) + (u - x_0)\varphi'_0(x_0)]^j \\
 &= \sum_{r=1}^j \binom{j}{r} [\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi'_0(x_0)]^r \\
 &\quad \times [\varphi'_0(x_0)]^{j-r} (u - x_0)^{j-r} \\
 &\quad + [\varphi'_0(x_0)]^j (u - x_0)^j.
 \end{aligned}$$

Hence, using (4.17) and (A3), we get on the set $\{u : |u - x_0| \leq Mn^{-1/(2k+1)}\}$

$$(\widehat{\varphi}_n(u) - \varphi_0(x_0))^j = o_p(n^{-k/(2k+1)})$$

for all $j \geq k + 1$.

In particular, this implies that

$$(4.21) \quad \widehat{\Psi}_{k,n,1}(u) = o_p(n^{-k/(2k+1)}),$$

uniformly in $u \in [x_0 - tn^{-1/(2k+1)}, x_0 + tn^{-1/(2k+1)}]$, where $|t| \leq M$, and

$$\begin{aligned}
 &\widehat{f}_n(u) - f_0(x_0) - f_0(x_0)(\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi'_0(x_0)) \\
 &\quad - f_0(x_0) \sum_{j=1}^k \frac{\varphi_0^{(j)}(x_0)}{j!} (u - x_0)^j = o_p(n^{-k/(2k+1)}).
 \end{aligned}$$

Using Lemma 4.2, the latter can be rewritten as

$$\widehat{f}_n(u) - f_0(x_0) - f_0(x_0)(\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi_0'(x_0)) - \sum_{j=1}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j - f_0(x_0) \frac{\varphi_0^{(k)}(x_0)}{k!} (u - x_0)^k = o_p(n^{-k/(2k+1)})$$

or, equivalently,

$$|\widehat{e}_n(x_0 + tn^{-1/(2k+1)}) - f_0(x_0)(\widehat{\varphi}_n(x_0 + tn^{-1/(2k+1)}) - \varphi_0(x_0) - n^{-1/(2k+1)}t\varphi_0'(x_0))| = o_p(n^{-k/(2k+1)})$$

uniformly in $|t| \leq M$. \square

THEOREM 4.6. *Let $K > 0$.*

(i) *If $\{Y_k(t), t \in \mathbb{R}\}$ is the canonical process defined in (2.1), then the localized process $\gamma_1 \mathbb{Y}_n^{\text{locmod}}(\gamma_2 \cdot)$ converges weakly in $C[-K, K]$ to Y_k , where*

$$(4.22) \quad \gamma_1 = \left(\frac{f_0(x_0)^{k-1} |\varphi_0^{(k)}(x_0)|^3}{[(k+2)!]^3} \right)^{1/(2k+1)},$$

$$(4.23) \quad \gamma_2 = \left(\frac{f_0(x_0) |\varphi_0^{(k)}(x_0)|^2}{[(k+2)!]^2} \right)^{1/(2k+1)}.$$

Equivalently, $\mathbb{Y}_n^{\text{locmod}}$ converges weakly in $C[-K, K]$ to the “driving process” $Y_{a,k,\sigma}$, where

$$(4.24) \quad Y_{k,a,\sigma}(t) := a \int_0^t W(s) ds - \sigma t^{k+2}$$

and where $a = 1/\sqrt{f_0(x_0)}$, $\sigma = |\varphi_0^{(k)}(x_0)|/(k+2)!$.

(ii) *The localized processes satisfy $\mathbb{Y}_n^{\text{locmod}}(t) - \widehat{H}_n^{\text{locmod}}(t) \geq 0$, for all $t \in \mathbb{R}$, with equality for all t such that $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{\mathcal{J}}_n(\widehat{\varphi}_n)$.*

(iii) *Both \widehat{A}_n and \widehat{B}_n defined above in (4.1) and (4.2) are tight.*

(iv) *The vector of processes*

$$(\widehat{H}_n^{\text{locmod}}, (\widehat{H}_n^{\text{locmod}})^{(1)}, (\widehat{H}_n^{\text{locmod}})^{(2)}, \mathbb{Y}_n^{\text{locmod}}, (\widehat{H}_n^{\text{locmod}})^{(3)}, (\mathbb{Y}_n^{\text{locmod}})^{(1)})$$

converges weakly in $(C[-K, K])^4 \times (D[-K, K])^2$, endowed with the product topology induced by the uniform topology on the spaces $C[-K, K]$ and the Skorohod topology on the spaces $D[-K, K]$ to the process

$$(H_{k,a,\sigma}, H_{k,a,\sigma}^{(1)}, H_{k,a,\sigma}^{(2)}, Y_{k,a,\sigma}, H_{k,a,\sigma}^{(3)}, Y_{k,a,\sigma}^{(1)}),$$

where $H_{k,a,\sigma}$ is the unique process on \mathbb{R} satisfying

$$(4.25) \quad \begin{cases} H_{k,a,\sigma}(t) \leq Y_{k,a,\sigma}(t), & \text{for all } t \in \mathbb{R}, \\ \int (H_{k,a,\sigma}(t) - Y_{k,a,\sigma}(t)) dH_{k,a,\sigma}^{(3)}(t) = 0, \\ H_{k,a,\sigma}^{(2)}, & \text{is concave.} \end{cases}$$

PROOF. (i) The first step will be to modify the local processes, that is, going from the “density” to the “log-density” level, in order to be able to exploit concavity of φ_0 and $\widehat{\varphi}_n$ and connect the local process to the limiting distribution obtained by Groeneboom, Jongbloed and Wellner (2001b) for estimating a convex density.

First, by Lemma 4.2, (4.19) and (A3), we can write

$$\begin{aligned} & f_0(x_0)^{-1} \left(\widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) \\ &= f_0(x_0)^{-1} \left(\widehat{f}_n(u) - f_0(x_0) - f_0(x_0) \sum_{j=1}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \right) \\ &= \widehat{\Psi}_{k,n,1}(u) + \sum_{j=1}^k \frac{1}{j!} [\widehat{\varphi}_n(u) - \varphi_0(x_0)]^j - \sum_{j=1}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \\ &= \widehat{\Psi}_{k,n,1}(u) + (\widehat{\varphi}_n(u) - \varphi_0(x_0) - \varphi_0'(x_0)(u - x_0)) \\ &\quad + \sum_{j=2}^k \frac{1}{j!} [\widehat{\varphi}_n(u) - \varphi_0(x_0)]^j - \sum_{j=2}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j \\ &=: (\widehat{\varphi}_n(u) - \varphi_0(x_0) - \varphi_0'(x_0)(u - x_0)) + \widehat{\Psi}_{k,n,2}(u), \end{aligned}$$

introducing the new remainder term

$$\begin{aligned} \widehat{\Psi}_{k,n,2}(u) &= \widehat{\Psi}_{k,n,1}(u) + \sum_{j=2}^k \frac{1}{j!} [\widehat{\varphi}_n(u) - \varphi_0(x_0)]^j \\ &\quad - \sum_{j=2}^{k-1} \frac{[\varphi_0'(x_0)]^j}{j!} (u - x_0)^j. \end{aligned} \tag{4.26}$$

Using (4.20) and (4.21) yields

$$\begin{aligned} & \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) \, du \, dv \\ &= t^2 n^{-2/(2k+1)} \sup_{u \in [x_0, v], v \in \mathcal{I}} |\widehat{\Psi}_{k,n,1}(u)| \\ &\quad + \sum_{j=2}^k \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0)]^j \, du \, dv \\ &\quad - \sum_{j=2}^{k-1} \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\varphi_0'(x_0)]^j (u - x_0)^j \, du \, dv \\ &= o_p(r_n^{-1}) \end{aligned}$$

$$\begin{aligned}
 & + \sum_{j=2}^k \frac{1}{j!} \sum_{l=1}^j \binom{j}{l} \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0) \\
 & \qquad \qquad \qquad - (u - x_0)\varphi'_0(x_0)]^l \\
 & \qquad \qquad \qquad \times (u - x_0)^{j-l} [\varphi'_0(x_0)]^{j-l} du dv \\
 & + \sum_{j=2}^k \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\varphi'_0(x_0)]^j (u - x_0)^j du dv \\
 & - \sum_{j=2}^{k-1} \frac{1}{j!} \int_{\mathcal{I}} \int_{x_0}^v [\varphi'_0(x_0)]^j (u - x_0)^j du dv \\
 & = o_p(r_n^{-1}) \\
 & + \sum_{j=2}^k \frac{1}{j!} \sum_{l=1}^j \binom{j}{l} \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0) \\
 & \qquad \qquad \qquad - (u - x_0)\varphi'_0(x_0)]^l \\
 & \qquad \qquad \qquad \times (u - x_0)^{j-l} [\varphi'_0(x_0)]^{j-l} du dv \\
 & + \frac{1}{k!} \int_{\mathcal{I}} \int_{x_0}^v (u - x_0)^k [\varphi'_0(x_0)]^k du dv.
 \end{aligned}$$

But by Lemma 4.5, one can easily show that, for $j = 2, \dots, k$ and $l = 1, \dots, j$,

$$\begin{aligned}
 & r_n \int_{\mathcal{I}} \int_{x_0}^v [\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi'_0(x_0)]^l (u - x_0)^{j-l} [\varphi'_0(x_0)]^{j-l} du dv \\
 & = O_p(n^{-[k(l-1)+(j-l)]/(2k+1)}) = o_p(1),
 \end{aligned}$$

uniformly in $|t| \leq M$. Similarly,

$$r_n \int_{\mathcal{I}} \int_{x_0}^v (u - x_0)^k [\varphi'_0(x_0)]^k du dv = \frac{[\varphi'_0(x_0)]^k}{(k+1)(k+2)} t^{k+2}.$$

Hence, it follows that

$$r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv = \frac{[\varphi'_0(x_0)]^k}{(k+2)!} t^{k+2} + o_p(1)$$

as $n \rightarrow \infty$, uniformly in $|t| \leq M$.

We turn now to the modified local processes, $\mathbb{Y}_n^{\text{locmod}}$ and $\widehat{H}_n^{\text{locmod}}$, defined in (4.3) and (4.4). It is not difficult to show that

$$(4.27) \quad \mathbb{Y}_n^{\text{locmod}}(t) = \frac{\mathbb{Y}_n^{\text{loc}}(t)}{f_0(x_0)} - r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) du dv$$

and

$$(4.28) \quad \widehat{H}_n^{\text{locmod}}(t) = \frac{\widehat{H}_n^{\text{loc}}(t)}{f_0(x_0)} - r_n \int_{\mathcal{I}} \int_{x_0}^v \widehat{\Psi}_{k,n,2}(u) \, du \, dv.$$

Note that the process $\widehat{H}_n^{\text{locmod}}$ is in fact similar to $\widehat{H}_n^{\text{loc}}$, except that it is defined in terms of the log-density φ_0 instead of the density f_0 . This can be more easily seen from its original expression given in (4.4). The second expression of $\widehat{H}_n^{\text{locmod}}$ given above is only useful for showing that it stays below $\mathbb{Y}_n^{\text{locmod}}$, while touching it at points t , such that $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{\mathcal{S}}_n(\widehat{\varphi}_n)$. The biggest advantage of considering this modified version is to be able to use concavity of φ_0 the same way [Groeneboom, Jongbloed and Wellner (2001b)] used convexity of the true estimated density g_0 . Their process $\widetilde{H}_n^{\text{loc}}$ resembles $\widehat{H}_n^{\text{locmod}}$ to a large extent (see page 1688), and by combining arguments similar to theirs with Lemma 4.2 and the results obtained above, it follows that

$$\begin{aligned} & \mathbb{Y}_n^{\text{locmod}}(t) \\ & \Rightarrow [f_0(x_0)]^{-1/2} \int_0^t W(s) \, ds + \frac{f_0^{(k)}(x_0)}{(k+2)!f_0(x_0)} t^{k+2} - \frac{[\varphi_0'(x_0)]^k}{(k+2)!} t^{k+2} \\ & = [f_0(x_0)]^{-1/2} \int_0^t W(s) \, ds + \frac{\varphi_0^{(k)}(x_0)}{(k+2)!} t^{k+2} \\ & = Y_{k,a,\sigma}(t) \quad \text{in } C[-K, K], \end{aligned}$$

where $a := [f_0(x_0)]^{-1/2}$, $\sigma := |\varphi_0^{(k)}(x_0)|/(k+2)!$, as in (4.24).

Now, let γ_1 and γ_2 be chosen, so that

$$\gamma_1 Y_{k,a,\sigma}(\gamma_2 t) \stackrel{d}{=} Y_k(t)$$

as processes where Y_k is the integrated Gaussian process defined in (2.1). Using the scaling property of Brownian motion [i.e., $\alpha^{-1/2}W(\alpha t) \stackrel{d}{=} W(t)$, for any $\alpha > 0$], we get

$$\gamma_1 \gamma_2^{3/2} = a^{-1} \quad \text{and} \quad \gamma_1 \gamma_2^{k+2} = \sigma^{-1}.$$

This yields γ_1 and γ_2 as given in (4.22) and (4.23), and hence,

$$\left(\begin{array}{c} n^{k/(2k+1)}(\widehat{\varphi}_n(x_0) - \varphi_0(x_0)) \\ n^{(k-1)/(2k+1)}(\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0)) \end{array} \right) \xrightarrow{d} f_0(x_0)^{-1} \left(\begin{array}{c} c_k(x_0, \varphi_0) H_k^{(2)}(0) \\ d_k(x_0, \varphi_0) H_k^{(3)}(0) \end{array} \right).$$

We get the explicit expression of the asymptotic constants $c_k(x_0, \varphi_0)$ and $d_k(x_0, \varphi_0)$ using the following relations:

$$(4.29) \quad f_0(x_0)^{-1} c_k(x_0, \varphi_0) = (\gamma_1 \gamma_2^2)^{-1} \quad \text{and}$$

$$(4.30) \quad f_0(x_0)^{-1} d_k(x_0, \varphi_0) = (\gamma_1 \gamma_2^3)^{-1}.$$

This is completely analogous to the derivations on page 1689 in Groeneboom, Jongbloed and Wellner (2001b), precisely

$$(4.31) \quad \begin{aligned} (\gamma_1 \widehat{H}_n^{\text{locmod}}(\gamma_2 t))^{(2)}(0) &= \gamma_1 \gamma_2^2 (\widehat{H}_n^{\text{locmod}})^{(2)}(0) \\ &= n^{k/(2k+1)} f_0(x_0) c_k(x_0, \varphi_0)^{-1} (\widehat{\varphi}_n(x_0) - \varphi_0(x_0)) \end{aligned}$$

and

$$(4.32) \quad \begin{aligned} (\gamma_1 \widehat{H}_n^{\text{locmod}}(\gamma_2 t))^{(3)}(0) &= \gamma_1 \gamma_2^3 (\widehat{H}_n^{\text{locmod}})^{(3)}(0) \\ &= n^{(k-1)/(2k+1)} f_0(x_0) d_k(x_0, \varphi_0)^{-1} (\widehat{\varphi}'_n(x_0) - \varphi'_0(x_0)). \end{aligned}$$

From (4.29) and (4.30), we get $c_k(x_0, \varphi_0)$ and $d_k(x_0, \varphi_0)$ as given in (2.2) and (2.3), and $C_k(x_0, \varphi_0)$ and $D_k(x_0, \varphi_0)$ as in (2.4) and (2.5).

(ii) Note that we can write

$$\mathbb{Y}_n^{\text{loc}}(t) - \widehat{H}_n^{\text{loc}}(t) = r_n (\mathbb{H}_n(x_n(t)) - \widehat{H}_n(x_n(t))) \geq 0$$

by making use of (1.1) and the specific choice of \widehat{A}_n and \widehat{B}_n . But, since we connect $\widehat{H}_n^{\text{locmod}}$ and $\mathbb{Y}_n^{\text{locmod}}$ to the “envelope,” the latter property needs primarily to hold for the modified processes. This can easily be established by considering (4.27) and (4.28), and hence it follows that

$$\mathbb{Y}_n^{\text{locmod}}(t) - \widehat{H}_n^{\text{locmod}}(t) \geq 0$$

for all $t \in \mathbb{R}$, with equality if $x_n(t) = x_0 + tn^{-1/(2k+1)} \in \widehat{\mathcal{S}}_n(\widehat{\varphi}_n)$.

(iii) To show that \widehat{A}_n and \widehat{B}_n are tight. By Theorem 4.3, we know that there exists $M > 0$ and $\tau \in \widehat{\mathcal{S}}(\widehat{\varphi}_n)$ such that $0 \leq x_0 - \tau \leq Mn^{-1/(2k+1)}$ with large probability. Now, using (4.11), we can write

$$\begin{aligned} |\widehat{A}_n| &\leq r_n s_n |(\widehat{F}_n(x_0) - \widehat{F}_n(\tau)) - (\mathbb{F}_n(x_0) - \mathbb{F}_n(\tau))| + r_n/n \\ &\leq r_n s_n \left| \int_{\tau}^{x_0} \left(\widehat{f}_n(u) - \sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j \right) du \right| \\ &\quad + r_n s_n \left| \int_{\tau}^{x_0} \left(\sum_{j=0}^{k-1} \frac{f_0^{(j)}(x_0)}{j!} (u - x_0)^j - f_0(u) \right) du \right| \\ &\quad + r_n s_n \left| \int_{\tau}^{x_0} d(\mathbb{F}_n - F_0) \right| + n^{-k/(2k+1)} \\ &:= \widehat{A}_{n1} + \widehat{A}_{n2} + \widehat{A}_{n3} + n^{-k/(2k+1)}. \end{aligned}$$

Now,

$$|\widehat{A}_{n1}| \leq r_n s_n \left| \int_{\tau}^{x_0} \widehat{e}_n(u) du - f_0(x_0) (\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0) \varphi'_0(x_0)) du \right|$$

$$\begin{aligned}
 & + r_n s_n f_0(x_0) \left| \int_{\tau}^{x_0} \left(\frac{[\varphi_0'(x_0)]^k}{k!} (u - x_0)^k \right) du \right| \\
 & + r_n s_n f_0(x_0) \left| \int_{\tau}^{x_0} (\widehat{\varphi}_n(u) - \varphi_0(x_0) - (u - x_0)\varphi_0'(x_0)) du \right| \\
 & \leq o_p(1) + O_p(r_n s_n (\tau - x_0)^{k+1}) + O_p(r_n s_n (\tau - x_0) n^{-k/(2k+1)}) \\
 & = O_p(1),
 \end{aligned}$$

where we used (4.18) and (4.17) to bound the first and last terms. To bound \widehat{A}_{n2} , we use Taylor approximation of $f_0(u)$ around x_0 to get

$$\begin{aligned}
 \widehat{A}_{n2} & \leq r_n \left| \int_{\tau}^{x_0} \frac{f_0^{(k)}(x_0)}{k!} (u - x_0)^k du \right| + r_n \left| \int_{\tau}^{x_0} (u - x_0)^k \varepsilon_n(u) du \right| \\
 & = O_p(1),
 \end{aligned}$$

where ε_n is a function such that $\|\varepsilon_n\| \rightarrow_p 0$ as $x_0 - \tau \rightarrow_p 0$. To bound \widehat{A}_{n3} , similar derivations as the ones used for bounding R_{2n} (see the proof of Lemma 4.4) can be employed where the perturbation function Δ_1 needs to be replaced by $\Delta_2(x) = 1_{[\tau, x_0]}(x)$.

At “one higher integration level,” similar computations can be used to show tightness of \widehat{B}_n .

(iv) The proof of this last part of the theorem is basically identical to that of Theorem 6.2 for the LSE in Groeneboom, Jongbloed and Wellner (2001b) and arguments similar to those of Groeneboom, Jongbloed and Wellner (2001a) or, alternatively, tightness plus uniqueness arguments along the lines of Groeneboom, Maathuis, and Wellner (2008). \square

PROOF OF THEOREM 2.1. The claimed joint convergence involving $\widehat{\varphi}_n$ and $\widehat{\varphi}'_n$ follows from part (iv) of Theorem 4.6 and the relations (4.31) and (4.32). The joint limiting distribution of $\widehat{f}_n(x_0) - f_0(x_0)$ and $\widehat{f}'_n(x_0) - f'_0(x_0)$ follows immediately by applying the delta-method. \square

4.3. Proofs for Section 3.

PROOF OF THEOREM 3.6. We first use the simple fact that \widehat{M}_n is the only point $x \in \mathbb{R}$ which satisfies

$$(4.33) \quad \widehat{\varphi}'_n(t) \begin{cases} > 0, & \text{if } t < x, \\ \leq 0, & \text{if } t \geq x. \end{cases}$$

This follows immediately from concavity of $\widehat{\varphi}_n$ and the definition of \widehat{M}_n . Note that $\widehat{\varphi}_n$ may have a flat region or “modal interval”; in this case, there exists an entire interval of points where the maximum is attained, and \widehat{M}_n is the left endpoint of this interval.

A tightness property of the process $H_2^{(3)}$, which follows from Lemma 2.7 of Groeneboom, Jongbloed and Wellner (2001b), is also needed to establish the limiting distribution of \widehat{M}_n : for any $\varepsilon > 0$ and $t \in \mathbb{R}$, there exists $C = C(\varepsilon)$ such that

$$P(|H_2^{(3)}(t) + 24t| > C) \leq \varepsilon.$$

In other words, one can view $H_2^{(3)}(t)$ as an “estimator” of the odd function $-24t$. Since C is independent of t , it follows that, for a fixed ε , $H_2^{(3)}(t) < 0$ [resp. $H_2^{(3)}(t) > 0$] for $t > 0$ (resp. $-t < 0$) big enough, with probability greater than $1 - \varepsilon$.

The sign of $H_2^{(3)}$ and uniqueness of \widehat{M}_n turn out to be crucial in determining the limiting distribution of the latter. From Theorem 4.6 and the two derivative relations, (4.31) and (4.32), it follows that

$$(4.34) \quad \begin{aligned} & \left(\begin{array}{l} n^{k/(2k+1)}(\widehat{\varphi}_n(x_0 + tn^{-1/(2k+1)}) - \varphi_0(x_0) - tn^{-1/(2k+1)}\varphi'_0(x_0)) \\ n^{(k-1)/(2k+1)}(\widehat{\varphi}'_n(x_0 + tn^{-1/(2k+1)}) - \varphi'_0(x_0)) \end{array} \right) \\ & \Rightarrow \left(\begin{array}{l} H_{k,a,\sigma}^{(2)}(t) \\ H_{k,a,\sigma}^{(3)}(t) \end{array} \right) \quad \text{in } C[-K, K] \times D[-K, K] \end{aligned}$$

for each $K > 0$, with the product topology induced by the uniform topology on $C[-K, K]$ and the Skorohod topology on $D[-K, K]$. Here, $H_{k,a,\sigma}$ is the unique process on \mathbb{R} satisfying (4.25). A similar result holds for the MLE of the log-concave density f_0 . When x_0 is replaced by the population mode $m_0 = M(f_0)$ and $k = 2$ the second weak convergence implies that

$$n^{1/5}(\widehat{\varphi}'_n(m_0 - Tn^{-1/5}) - \varphi'_0(m_0)) \xrightarrow{d} H_{2,a,\sigma}^{(3)}(-T)$$

and

$$n^{1/5}(\widehat{\varphi}'_n(m_0 + Tn^{-1/5}) - \varphi'_0(m_0)) \xrightarrow{d} H_{2,a,\sigma}^{(3)}(T).$$

For $T > 0$ large enough, this in turn implies that, for $\varepsilon > 0$, we can find $N \in \mathbb{N} \setminus \{0\}$ such that, for all $n > N$, we have

$$P(\widehat{\varphi}'_n(m_0 - Tn^{-1/5}) > 0 \text{ and } \widehat{\varphi}'_n(m_0 + Tn^{-1/5}) < 0) > 1 - \varepsilon.$$

Using the property of \widehat{M}_n in (4.33), it follows that

$$P(\widehat{M}_n \in [m_0 - Tn^{-1/5}, m_0 + Tn^{-1/5}]) > 1 - \varepsilon$$

for all $n > N$.

We first conclude that $\widehat{M}_n - m_0 = O_p(n^{-1/5})$. Then, we note that

$$n^{1/5}(\widehat{M}_n - m_0) = M(\mathbb{Z}_n),$$

where

$$\begin{aligned} \mathbb{Z}_n(t) &= n^{2/5}(\widehat{\varphi}_n(m_0 + tn^{-1/5}) - \varphi_0(m_0)) \\ &\Rightarrow \mathbb{Z}(t) := H_{2,a,\sigma}^{(2)}(t) \quad \text{in } C([-K, K]) \end{aligned}$$

for each $K > 0$, by (4.34) with $k = 2$. Thus, by the argmax continuous mapping theorem [see, e.g., van der Vaart and Wellner (1996), page 286] it follows that

$$M(\mathbb{Z}_n) \xrightarrow{d} M(\mathbb{Z}) = M(H_{2,a,\sigma}^{(2)}),$$

where $\mathbb{Z} = H_{2,a,\sigma}^{(2)}$, $a = 1/\sqrt{f_0(m_0)}$, and $\sigma = |\varphi_0^{(2)}(m_0)|/4!$.

Note that $H_{2,a,\sigma}$ is related to the “driving process” $Y_{2,a,\sigma}$ with $a = 1/\sqrt{f_0(m_0)}$, $\sigma = |\varphi_0^{(2)}(m_0)|/4!$ as in (4.24) with $k = 2$. Now, $\gamma_1 Y_{2,a,\sigma}(\gamma_2 t) \stackrel{d}{=} Y_2(t)$ as processes where $Y_2 := Y_{2,1,1}$. Thus, it also holds that

$$\gamma_1 H_{2,a,\sigma}(\gamma_2 t) \stackrel{d}{=} H_2(t) \quad \text{and} \quad \gamma_1 \gamma_2^2 H_{2,a,\sigma}^{(2)}(\gamma_2 t) \stackrel{d}{=} H_2^{(2)}(t),$$

or, equivalently, $H_{2,a,\sigma}^{(2)}(v) \stackrel{d}{=} H_2^{(2)}(v/\gamma_2)/(\gamma_1 \gamma_2^2)$. Since $M(dg(c \cdot)) = c^{-1}M(g)$ for $c, d > 0$, it follows that

$$M(H_{2,a,\sigma}^{(2)}) \stackrel{d}{=} M\left(\frac{1}{\gamma_1 \gamma_2^2} H_2^{(2)}(\cdot/\gamma_2)\right) \stackrel{d}{=} \gamma_2 M(H_2^{(2)}),$$

where

$$\gamma_2 = \left(f_0(m_0) \frac{|\varphi_0^{(2)}(m_0)|^2}{(4!)^2}\right)^{-1/5} = \left(\frac{(4!)^2 f_0(m_0)}{f_0''(m_0)^2}\right)^{1/5}$$

by direct computation using $f_0'(m_0) = 0 = \varphi_0'(m_0)$ and Lemma 4.2. \square

Acknowledgments. This research was initiated while Kaspar Rufibach was visiting the Institute for Mathematical Stochastics at the University of Göttingen, Germany and while Fadoua Balabdaoui was visiting the Institute of Mathematical Statistics and Actuarial Science at the University of Bern, Switzerland. We would like to thank both institutions for their hospitality. We also thank two referees and an Associate Editor for several suggestions leading to improvements of the exposition.

REFERENCES

- AN, M. Y. (1998). Logconcavity versus logconvexity: A complete characterization. *J. Econom. Theory* **80** 350–369. [MR1637480](#)
- ANEVSKI, D. and HÖSSJER, O. (2006). A general asymptotic scheme for inference under order restrictions. *Ann. Statist.* **34** 1874–1930. [MR2283721](#)
- BAGNOLI, M. and BERGSTROM, T. (2005). Log-concave probability and its applications. *Econom. Theory* **26** 445–469. [MR2213177](#)
- BALABDAOUI, F. and WELLNER, J. A. (2007). Estimation of a k -monotone density: Limiting distribution theory and the spline connection. *Ann. Statist.* **35** 2536–2564. [MR2382657](#)
- BARLOW, R. E. and PROSCHAN, F. (1975). *Statistical Theory of Reliability and Life Testing*. Holt, Rinehart and Winston, New York. [MR0438625](#)
- BIRGÉ, L. (1989). The Grenander estimator: A nonasymptotic approach. *Ann. Statist.* **17** 1532–1549. [MR1026298](#)

- BIRGÉ, L. (1997). Estimation of unimodal densities without smoothness assumptions. *Ann. Statist.* **25** 970–981. [MR1447736](#)
- BRUNNER, L. J. and LO, A. Y. (1989). Bayes methods for a symmetric unimodal density and its mode. *Ann. Statist.* **17** 1550–1566. [MR1026299](#)
- CHANG, G. and WALTHER, G. (2007). Clustering with mixtures of log-concave distributions. *Comput. Statist. Data Anal.* **51** 6242–6251. [MR2408591](#)
- CHERNOFF, H. (1964). Estimation of the mode. *Ann. Inst. Statist. Math.* **16** 31–41. [MR0172382](#)
- CULE, M., GRAMACY, R. and SAMWORTH, R. (2007). LogConcDEAD: Maximum likelihood estimation of a log-concave density. R package version 1.0-5.
- DALENIUS, T. (1965). The mode—a neglected statistical parameter. *J. Roy. Statist. Soc. Ser. A* **128** 110–117. [MR0185720](#)
- DHARMADHIKARI, S. and JOAG-DEV, K. (1988). *Unimodality, Convexity, and Applications*. Academic Press, Boston, MA. [MR0954608](#)
- DÜMBGEN, L., HÜSLER, A. and RUFIBACH, K. (2007). Active set and EM algorithms for log-concave densities based on complete and censored data. Technical report, Univ. Bern. Available at [arXiv:0707.4643](#).
- DÜMBGEN, L. and RUFIBACH, K. (2009). Maximum likelihood estimation of a log-concave density and its distribution function: Basic properties and uniform consistency. *Bernoulli* **15**. To appear. Available at [arXiv:0709.0334](#).
- EDDY, W. F. (1980). Optimum kernel estimators of the mode. *Ann. Statist.* **8** 870–882. [MR0572631](#)
- EDDY, W. F. (1982). The asymptotic distributions of kernel estimators of the mode. *Z. Wahrsch. Verw. Gebiete* **59** 279–290. [MR0721626](#)
- EKBLOM, H. (1972). A Monte Carlo investigation of mode estimators in small samples. *Appl. Statist.* **21** 177–184.
- GRENANDER, U. (1956). On the theory of mortality measurement. II. *Skand. Aktuarietidskr.* **39** 125–153 (1957). [MR0093415](#)
- GRENANDER, U. (1965). Some direct estimates of the mode. *Ann. Math. Statist.* **36** 131–138. [MR0170409](#)
- GROENEBOOM, P. (1996). Lectures on inverse problems. In *Lectures on Probability Theory and Statistics (Saint-Flour, 1994)*. *Lecture Notes in Math.* **1648** 67–164. Springer, Berlin. [MR1600884](#)
- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2001a). A canonical process for estimation of convex functions: The “invelope” of integrated Brownian motion $+t^4$. *Ann. Statist.* **29** 1620–1652. [MR1891741](#)
- GROENEBOOM, P., JONGBLOED, G. and WELLNER, J. A. (2001b). Estimation of a convex function: Characterizations and asymptotic theory. *Ann. Statist.* **29** 1653–1698. [MR1891742](#)
- GROENEBOOM, P., MAATHUIS, M. H. and WELLNER, J. A. (2008). Current status data with competing risks: Limiting distribution of the MLE. *Ann. Statist.* **36** 1064–1089. [MR2418649](#)
- GRUND, B. and HALL, P. (1995). On the minimisation of L^p error in mode estimation. *Ann. Statist.* **23** 2264–2284. [MR1389874](#)
- HALL, P. (1982). Asymptotic theory of Grenander’s mode estimator. *Z. Wahrsch. Verw. Gebiete* **60** 315–334. [MR0664420](#)
- HAS’MINSKIĬ, R. Z. (1979). Lower bound for the risks of nonparametric estimates of the mode. In *Contributions to Statistics* 91–97. Reidel, Dordrecht. [MR0561262](#)
- HERRMANN, E. and ZIEGLER, K. (2004). Rates on consistency for nonparametric estimation of the mode in absence of smoothness assumptions. *Statist. Probab. Lett.* **68** 359–368. [MR2081114](#)
- HO, M.-W. (2006a). Bayes estimation of a symmetric unimodal density via S-paths. *J. Comput. Graph. Statist.* **15** 848–860. [MR2273482](#)
- HO, M.-W. (2006b). Bayes estimation of a unimodal density via S-paths. Technical report, National University of Singapore. Available at <http://arxiv.org/math/0609056>.
- IBRAGIMOV, I. A. (1956). On the composition of unimodal distributions. *Teor. Veroyatnost. i Primenen.* **1** 283–288. [MR0087249](#)

- JANKOWSKI, H. and WELLNER, J. A. (2007). Nonparametric estimation of a convex bathtub-shaped hazard function. Technical report, Univ. Washington.
- JONGBLOED, G. (1995). Three statistical inverse problems. Ph.D. thesis, Delft Univ. Technology.
- JONGBLOED, G. (2000). Minimax lower bounds and moduli of continuity. *Statist. Probab. Lett.* **50** 279–284. [MR1792307](#)
- KIM, J. and POLLARD, D. (1990). Cube root asymptotics. *Ann. Statist.* **18** 191–219. [MR1041391](#)
- KLEMELÄ, J. (2005). Adaptive estimation of the mode of a multivariate density. *J. Nonparametr. Statist.* **17** 83–105. [MR2112688](#)
- LEPSKIĪ, O. V. (1991). Asymptotically minimax adaptive estimation. I. Upper bounds. Optimally adaptive estimates. *Teor. Veroyatnost. i Primenen.* **36** 645–659. [MR1147167](#)
- LEPSKIĪ, O. V. (1992). Asymptotically minimax adaptive estimation. II. Schemes without optimal adaptation. Adaptive estimates. *Teor. Veroyatnost. i Primenen.* **37** 468–481. [MR1214353](#)
- LEURGANS, S. (1982). Asymptotic distributions of slope-of-greatest-convex-minorant estimators. *Ann. Statist.* **10** 287–296. [MR0642740](#)
- MARSHALL, A. W. and OLKIN, I. (1979). *Inequalities: Theory of Majorization and Its Applications. Mathematics in Science and Engineering* **143**. Academic Press [Harcourt Brace Jovanovich Publishers], New York. [MR0552278](#)
- MARSHALL, A. W. and OLKIN, I. (2007). *Life Distributions*. Springer, New York. [MR2344835](#)
- MEYER, M. C. (2001). An alternative unimodal density estimator with a consistent estimate of the mode. *Statist. Sinica* **11** 1159–1174. [MR1867337](#)
- MEYER, M. C. and WOODROOFE, M. (2004). Consistent maximum likelihood estimation of a unimodal density using shape restrictions. *Canad. J. Statist.* **32** 85–100. [MR2060547](#)
- MÜLLER, H.-G. (1989). Adaptive nonparametric peak estimation. *Ann. Statist.* **17** 1053–1069. [MR1015137](#)
- MÜLLER, S. and RUFIBACH, K. (2009). Smoothed tail index estimation. *J. Statist. Comput. Simulation*. To appear. Available at [arXiv:math/0612140](#).
- PAL, J. K., WOODROOFE, M. B. and MEYER, M. C. (2007). Estimating a Polya frequency function. In *Complex Datasets and Inverse Problems: Tomography, Networks, and Beyond. IMS Lecture Notes-Monograph Series* **54** 239–249.
- PARZEN, E. (1962). On estimation of a probability density function and mode. *Ann. Math. Statist.* **33** 1065–1076. [MR0143282](#)
- PRAKASA RAO, B. L. S. (1969). Estimation of a unimodal density. *Sankhyā Ser. A* **31** 23–36. [MR0267677](#)
- ROMANO, J. P. (1988). On weak convergence and optimality of kernel density estimates of the mode. *Ann. Statist.* **16** 629–647. [MR0947566](#)
- RUFIBACH, K. (2006). Log-concave density estimation and bump hunting for I.I.D. observations. Ph.D. thesis, Univ. Bern and Göttingen.
- RUFIBACH, K. (2007). Computing maximum likelihood estimators of a log-concave density function. *J. Statist. Comput. Simulation* **77** 561–574. [MR2407642](#)
- RUFIBACH, K. and DÜMBGEN, L. (2007). Logcondens: estimate a log-concave probability density from i.i.d. observations. R package version 1.3.1.
- VAN DER VAART, A. W. and WELLNER, J. A. (1996). *Weak Convergence and Empirical Processes with Applications in Statistics*. Springer, New York. [MR1385671](#)
- VENTER, J. H. (1967). On estimation of the mode. *Ann. Math. Statist* **38** 1446–1455. [MR0216698](#)
- VIEU, P. (1996). A note on density mode estimation. *Statist. Probab. Lett.* **26** 297–307. [MR1393913](#)
- WALTHER, G. (2002). Detecting the presence of mixing with multiscale maximum likelihood. *J. Amer. Statist. Assoc.* **97** 508–513. [MR1941467](#)
- WEGMAN, E. J. (1970a). Maximum likelihood estimation of a unimodal density function. *Ann. Math. Statist.* **41** 457–471. [MR0254995](#)
- WEGMAN, E. J. (1970b). Maximum likelihood estimation of a unimodal density. II. *Ann. Math. Statist.* **41** 2169–2174. [MR0267681](#)

- WEGMAN, E. J. (1971). A note on the estimation of the mode. *Ann. Math. Statist.* **42** 1909–1915. [MR0297072](#)
- WRIGHT, F. T. (1981). The asymptotic behavior of monotone regression estimates. *Ann. Statist.* **9** 443–448. [MR0606630](#)
- ZIEGLER, K. (2004). Adaptive kernel estimation of the mode in a nonparametric random design regression model. *Probab. Math. Statist.* **24** 213–235. [MR2157203](#)

F. BALABDAOUI
CENTRE DE RECHERCHE EN MATHÉMATIQUES DE LA DÉCISION
UNIVERSITÉ PARIS-DAUPHINE
PARIS
FRANCE
E-MAIL: fadoua@ceremade.dauphine.fr

K. RUFIBACH
INSTITUTE OF SOCIAL
AND PREVENTIVE MEDICINE
BIostatistics UNIT
UNIVERSITY OF ZÜRICH
CH-8001 ZÜRICH
SWITZERLAND
E-MAIL: kaspar.rufibach@ifspm.uzh.ch

J. A. WELLNER
DEPARTMENT OF STATISTICS
UNIVERSITY OF WASHINGTON
BOX 354322
SEATTLE, WASHINGTON 98195-4322
USA
E-MAIL: jaw@stat.washington.edu