

Bi- s^* -Concave Distributions



Jon A. Wellner (Seattle)

Non- and Semiparametric Statistics

*In Honor of Arnold Janssen,
On the occasion of his 65th Birthday*

Non- and Semiparametric Statistics:

**In Honor of Arnold Janssen
Dusseldorf, Germany**

July 21-22, 2017

Based on joint work with: [Nilanjana Laha](#)

Outline

- 0. A mysterious condition: quantile process theory
- 1. Bi-log-concavity.
- 2. Questions and examples.
- 3. Bi- s^* -concavity.
- 4. Open questions.

1. A Mysterious condition: quantile process theory

- Let X_1, \dots, X_n be i.i.d. F ,
absolutely continuous with density f .
- Let \mathbb{F}_n denote the empirical distribution function of the X_i 's:
 $\mathbb{F}_n(x) = n^{-1} \sum_{i=1}^n \mathbf{1}\{X_i \leq x\}$.
- Let \mathbb{F}_n^{-1} denote the empirical quantile function, and let F^{-1} denote the population quantile function,
where $F^{-1}(t) \equiv \inf\{x : F(x) \geq t\}$, $0 < t < 1$.

- The standardized quantile process \mathbb{Q}_n is defined by

$$\mathbb{Q}_n(t) \equiv g(t) \sqrt{n} (\mathbb{F}_n^{-1}(t) - F^{-1}(t)) \quad \text{for } 0 < t < 1.$$

where

$$g(t) \equiv f(F^{-1}(t)).$$

is the *density quantile function*.

Now suppose that $X_i \equiv F^{-1}(\xi_i)$ for $1 \leq i \leq n$ where:

- ξ_1, \dots, ξ_n are i.i.d. Uniform(0, 1) random variables.
- \mathbb{G}_n is the empirical d.f. of the ξ_i 's.
- \mathbb{G}_n^{-1} is the empirical quantile function of the ξ_i 's.
- $\mathbb{V}_n(t) \equiv \sqrt{n}(\mathbb{G}_n^{-1}(t) - t)$ is the *uniform quantile process*.

Csörgő and Révész (1978) imposed the following mysterious condition in their study of the asymptotic equivalence of \mathbb{V}_n and \mathbb{Q}_n^0 , the version of \mathbb{Q}_n with the X_i 's constructed in terms of the ξ_i 's as above.

Suppose

$$\gamma(F) \equiv \sup_{x \in J(F)} F(x)(1 - F(x)) \frac{|f'(x)|}{f^2(x)} \leq \text{some } M < \infty. \quad (1)$$

Then, for any (small) $r > 0$

$$\|\mathbb{Q}_n^0 - \mathbb{V}_n\|_\infty = O\left(n^{-1/2}(\log\log n)^M (\log n)^{(1+r)(M-1)}\right) \text{ a.s.}$$

Define the $CR(x)$ and $CR_m(x)$ functions as follows:

$$CR(x) \equiv F(x)(1 - F(x)) \frac{|f'(x)|}{f^2(x)},$$
$$CR_m(x) \equiv \min\{F(x), 1 - F(x)\} \frac{|f'(x)|}{f^2(x)}.$$

The condition (1) has also appeared in the study of transportation distances between the empirical measure and true measure \mathbb{P}_n and P on \mathbb{R} ; see e.g.

- del Barrio, E., Giné, E., and Utzet, F. (2005). Asymptotics for L2 functionals of the empirical quantile process, with applications to tests of fit based on weighted Wasserstein distances. *Bernoulli* **11**, 131 - 189.
- Bobkov, S. and Ledoux, M. (2014). One - Dimensional empirical measures, order statistics, and Kantorovich transport distances. *Memoirs of the American Mathematical Society*, to appear. (especially see p. 45)

2. Bi-log-concavity


Definition: Dümbgen, Kolesnyk, and Wilke (2017)

A distribution function F on \mathbb{R} is **bi-log-concave** if both $\log F$ and $\log(1 - F)$ are concave functions from \mathbb{R} to $[-\infty, 0]$.


- DKW (2017) noted that if F has log-concave density $f = F'$, then F is bi-log-concave.
- But ... bi-log-concavity of F is a much weaker constraint:
 - ▶ While any log-concave density is unimodal,
 - ▶ a bi-log-concave distribution function F may have a density with an arbitrary number of modes; e.g.

$$f_{k,a}(x) = \left(1 + \frac{a \sin(2\pi kx)}{k\pi}\right) 1_{[0,1]}(x) / C(k, a).$$


for a small, is bi-log-concave




Gaussian density ϕ



$CR(x)$ and $CR_m(x)$ for $f = \phi$




Mixed Gaussian density,
 $2^{-1}\phi(x + \delta) + 2^{-1}\phi(x - \delta)$, $\delta = 1.37$




CR functions, mixed Gaussian density:

$$2^{-1}\phi(x + \delta) + 2^{-1}\phi(x - \delta), \delta = 1.37$$



Multi-modal perturbed uniform density, $f_{k,a}$ with $k = 4$, $a = 0.795$



C-R functions, multi-modal perturbed uniform density:

$$f_{k,a} \text{ with } k = 4, a = 0.0795$$

Let

$$J(F) \equiv \{x \in \mathbb{R} : 0 < F(x) < 1\}.$$

Then a distribution function F is non-degenerate if $J(F) \neq \emptyset$.

Theorem 1. (DKW-2017).

If F is non-degenerate, the following four statements are equivalent:

(i) F is [bi-log-concave](#).

(ii) F is continuous on \mathbb{R} and differentiable on $J(F)$ with derivative $f = F'$ such that

$$F(x+t) \begin{cases} \leq F(x) \exp\left(\frac{f(x)}{F(x)}t\right), \\ \geq 1 - (1 - F(x)) \exp\left(-\frac{f(x)}{1-F(x)}t\right) \end{cases}$$

for all $x \in J(F)$ and $t \in \mathbb{R}$.

(iii) F is continuous on \mathbb{R} and differentiable on $J(F)$ with derivative $f = F'$ such that the hazard function $f/(1 - F)$ is non-decreasing and the reverse hazard function f/F is non-increasing on $J(F)$.

(iv) F is continuous on \mathbb{R} and differentiable on $J(F)$ with bounded and strictly positive derivative $f = F'$. Furthermore, f is locally Lipschitz continuous on $J(F)$ with L^1 -derivative $f' = F''$ satisfying

$$\frac{-f^2}{1 - F} \leq f' \leq \frac{f^2}{F},$$

or, equivalently,

$$\frac{-f}{1 - F} \leq \frac{f'}{f} \leq \frac{f}{F}.$$

Corollary. If F is bi-log-concave

$$\tilde{\gamma}(F) \equiv \sup_{x \in \mathbb{R}} \{F(x) \wedge (1 - F(x))\} \frac{|f'(x)|}{f^2(x)} \leq 1,$$

$$\gamma(F) \equiv \sup_{x \in \mathbb{R}} F(x)(1 - F(x)) \frac{|f'(x)|}{f^2(x)} \leq 1.$$

3. Questions and some examples

If a density f on \mathbb{R}^d is of the form

$$f(x) \equiv f_\varphi(x) = \begin{cases} (\varphi(x))^{1/s}, & \varphi \text{ convex, if } s < 0 \\ \exp(-\varphi(x)), & \varphi \text{ convex, if } s = 0 \\ (\varphi(x))^{1/s}, & \varphi \text{ concave, if } s > 0, \end{cases}$$

then f is **s -concave**.

The classes of all densities f on \mathbb{R}^d of these forms are called the classes of s -concave densities, \mathcal{P}_s . The following inclusions hold: if $-\infty < s < 0 < r < \infty$, then

$$\mathcal{P}_\infty \subset \mathcal{P}_r \subset \mathcal{P}_0 \subset \mathcal{P}_s \subset \mathcal{P}_{-\infty}$$

Questions:

- **Q1:** What if the density f is s -concave with $s \neq 0$. In particular, what if $f \in \mathcal{P}_s$ with $s < 0$ where we know (Borell, Brascamp & Lieb, Rinott, ...)

$$\mathcal{P}_{-\infty} \supset \mathcal{P}_s \supset \mathcal{P}_0 \supset \mathcal{P}_r \supset \mathcal{P}_{\infty}$$

for $-\infty < s < 0 < r < \infty$?

- **Q2:** If $f \in \mathcal{P}_s$, is there a class of bi- s^* -concave distribution functions F with the property that F and $1 - F$ are s^* -concave?
- **Q3:** Is there an analogue of Theorem 1 including an analogue of Theorem 1(iv) with the corollary that $\gamma(F)$ is bounded by some function of s ?

From Borell, Brascamp, & Lieb, Rinott, we know that if $f \in \mathcal{P}_s$ for $s > -1$, then the measure $P_f(A) = \int_A f d\lambda$ for Borel sets A is t -concave with $t = s/(1+s) \equiv s^*$ for $s > -1$. Thus by taking $A = (-\infty, x]$, it follows that $x \mapsto F(x)$ is s^* -concave; similarly, taking $A = [x, \infty)$ it follows that $x \mapsto 1 - F(x)$ is s^* -concave.

Example 1. t_r densities: $s = -1/(1+r)$; $s^* = s/(1+s) = -1/r$. Suppose that

$$f_r(x) = \frac{C_r}{\left(1 + \frac{x^2}{r}\right)^{(r+1)/2}}$$


where $C_r = \Gamma((r+1)/2)/(\sqrt{\pi}\Gamma(r/2))$. Then f_r is s -concave for all $s \leq -(1+r)^{-1}$. By the Borell-Brascamp-Lieb-Rinott correspondence between s -concave densities we know that


$$x \mapsto F_r(x)^{s^*} \quad \text{and} \quad x \mapsto (1 - F_r(x))^{s^*}$$


are convex.


Here are some plots, for $r \in \{1/8, 1/4, 1/2, 1, 4, 16\}$,
and hence $s \in \{-8/9, -4/5, -2/3, -1/2, -1/3, -1/5\}$:

- f_r ,
- f_r^s , $s = -1/(1 + r)$.
- $f_r/(1 - F_r)^{1-s^*}$, $s^* = s/(1 + s) = -1/r$.
- $CRm(x, f) \equiv \min\{F(x), 1 - F(x)\}f'(x)/f(x)^2$ for $f = f_r$.
 $CR(x, f) \equiv F(x)(1 - F(x))f'(x)/f(x)^2$.










Here the black bounding lines at the top and bottom are given by


$$1 - s^* = \frac{1}{1 + s} = \frac{1}{1 - 8/9} = 9 \quad \text{since} \quad s = -\frac{1}{1 + 1/8} = -\frac{8}{9}.$$


Example 2. (Mixtures of t_r) Suppose that

$$f(x) = f(x; r, \delta) \equiv \frac{1}{2}g_r(x - \delta) + \frac{1}{2}g_r(x + \delta)$$

where g_r is the t_r -density in Example 1 and where $\delta > 0$ is not too large. For example here are Figures 2 - 5 of Laha & W (2017). Showing g_r with $r = 1$ and $\delta = 1.3$








Example 3. (symmetric beta densities). Now consider the family of s -concave densities with $s > 0$ given for any $r \in (0, \infty)$ by


$$f_r(x) = \sqrt{r}C_r(1 - x^2)^{r/2}1_{[-1,1]}(x)$$

where $C_r \equiv \Gamma((3 + r)/2)/(\sqrt{\pi r}\Gamma(1 + r/2))$. Then $f_r \in \mathcal{P}_s$. with $s = 2/r$. Here are some plots, for $r \in \{1/8, 1/2, 2, 4, 8, 16\}$, and hence $s \in \{16, 4, 1, 1/2, 1/4, 1/8\}$:


- f_r ,
- f_r^s , $s = 2/r$.
- $f_r/(1 - F_r)^{1-s^*}$, $s^* = s/(1 + s) =$.
- $CRm(x, f) \equiv \min\{F(x), 1 - F(x)\}f'(x)/f(x)^2$ for $f = f_r$.
 $CR(x, f) \equiv F(x)(1 - F(x))f'(x)/f(x)^2$.




Symmetrized beta densities f_r with $r \in \{1/8, 1/4, 1/2, 2, 4, 16\}$



Powers of symmetrized beta densities $f_r^s = f_r^{2/r}$
with $r \in \{1/8, 1/4, 1/2, 2, 4, 16\}$





$CRm(x)$ for f_r symmetrized Beta, $r \in \{1/8, 1/4, 1/2, 2, 4, 16\}$
 Here the black bounding lines at the top and bottom are given by the bound for the biggest class, namely for $r = 16$, so $s = 1/8$ and

$$1 - s^* = \frac{1}{1 + s} = \frac{1}{1 + 1/8} = \frac{8}{9} \quad \text{since} \quad s = \frac{2}{16} = \frac{1}{8}.$$

3. Bi- s^* -concave distributions

Definition.

- For $s \in (-1, \infty)$, let $s^* \equiv s/(1 + s) \in (-\infty, 1]$.
- For $s \in (-1, 0)$, a distribution function F on \mathbb{R} is **bi- s^* -concave** if both $x \mapsto F^{s^*}(x)$ and $x \mapsto (1 - F)^{s^*}(x)$ are **convex** functions of $x \in J(F)$.
- For $s \in (0, \infty)$, F on \mathbb{R} is **bi- s^* -concave** if $x \mapsto F^{s^*}(x)$ is concave for $x \in (\inf J(F), \infty)$ and $x \mapsto (1 - F)^{s^*}(x)$ is concave for $x \in (-\infty, \sup J(F))$.
- For $s = 0$, F on \mathbb{R} is **bi-0-concave** or **bi-log-concave** if both $x \mapsto \log F(x)$ and $x \mapsto \log(1 - F(x))$ are **concave** functions of $x \in J(F)$.

Theorem 2. (Bi- s^* -characterization theorem) Let $s \in (-1, \infty]$. For a non-degenerate distribution function F the following four statements are equivalent:

(i) F is bi- s^* -concave.

(ii) F is continuous on \mathbb{R} and differentiable on $J(F)$ with derivative $f = F'$. Moreover when $s \leq 0$,

$$F(x+t) \begin{cases} \leq F(x) \cdot \left(1 + s^* \frac{f(x)}{F(x)} t\right)_+^{1/s^*} \\ \geq 1 - (1 - F(x)) \cdot \left(1 - s^* \frac{f(x)}{1-F(x)} t\right)_+^{1/s^*} \end{cases} \quad (2)$$

for all $x \in \mathbb{R}$ and $t \in \mathbb{R}$. When $s > 0$,

$$F(x+t) \begin{cases} \leq F(x) \cdot \left(1 + s^* \frac{f(x)}{F(x)} t\right)_+^{1/s^*}, & \text{for } t \in (a-x, \infty) \\ \geq 1 - (1 - F(x)) \cdot \left(1 - s^* \frac{f(x)}{1-F(x)} t\right)_+^{1/s^*}, & \text{for } t \in (-\infty, b-x) \end{cases} \quad (3)$$

for all $x \in J(F)$.

(iii) F is continuous on \mathbb{R} and differentiable on $J(F)$ with derivative $f = F'$ such that the s^* -hazard function $f/(1-F)^{1-s^*}$ is non-decreasing, and the reverse s^* -hazard function f/F^{1-s^*} is non-increasing on $J(F)$.

(iv) F is continuous on \mathbb{R} and differentiable on $J(F)$ with bounded and strictly positive derivative $f = F'$. Furthermore, f is locally Lipschitz-continuous on $J(F)$ with L^1 -derivative $f' = F''$ satisfying

$$-(1-s^*)\frac{f^2}{1-F} \leq f' \leq (1-s^*)\frac{f^2}{F}. \quad (4)$$

Corollary.

Suppose that F is bi- s^* -concave for $s \in (-1, \infty]$. Then

$$\gamma(F) = \sup_{x \in J(F)} F(x)(1 - F(x)) \frac{|f'(x)|}{f^2(x)} \leq 1 - s^* = \frac{1}{1 + s},$$

and

$$\tilde{\gamma}(F) = \sup_{x \in J(F)} \min\{F(x), 1 - F(x)\} \frac{|f'(x)|}{f^2(x)} \leq 1 - s^* = \frac{1}{1 + s}.$$

Questions and further problems:

- Q1.** Application of bi- s^* -concavity to construction of confidence bands for F . For $s = 0$, this has been implemented by DKW (2017).
- Q2.** Can anything be said when f is s -concave with $s \leq -1$?
- Q3.** Bi-log-concave or bi- s^* -concave in higher dimensions?
- Q4.** What are the “right” hypotheses for the study of transportation (Wasserstein) distances for empirical measures on \mathbb{R}^d with $d \geq 2$?

Selected references:

- Dümbgen, L., Kolesnyk, P., and Wilke, R. (2017). Bi-log-concave distribution functions. *J. Statist. Planning and Inference* **184**, 1 - 17.
- del Barrio, E., Giné, E., and Utzet, F. (2005). Asymptotics for L_2 functionals of the empirical quantile process, with applications to tests of fit based on weighted Wasserstein distances. *Bernoulli* **11**, 131 - 189.
- Laha, N. and Wellner, J. A. (2017). Bi- s^* -concave distributions. Submitted. Available as [arXiv:1705.00252](https://arxiv.org/abs/1705.00252).

Fröhlichen Geburtstag Arnold!